



Cambridge
Elements

Quantitative and
Computational Methods
for the Social Sciences

Quantile Modeling for Political Research

Xiao Lu

ISSN 2398-4023 (online)

ISSN 2514-3794 (print)

Cambridge Elements[≡]

Elements in Quantitative and Computational
Methods for the Social Sciences

edited by

R. Michael Alvarez

California Institute of Technology

Betsy Sinclair

Washington University in St. Louis

QUANTILE MODELING FOR POLITICAL RESEARCH

Xiao Lu

Peking University



CAMBRIDGE
UNIVERSITY PRESS



Shaftesbury Road, Cambridge CB2 8EA, United Kingdom
One Liberty Plaza, 20th Floor, New York, NY 10006, USA
477 Williamstown Road, Port Melbourne, VIC 3207, Australia
314–321, 3rd Floor, Plot 3, Splendor Forum, Jasola District Centre,
New Delhi – 110025, India

Cambridge University Press is part of Cambridge University Press & Assessment,
a department of the University of Cambridge.

We share the University's mission to contribute to society through the pursuit of
education, learning and research at the highest international levels of excellence.

www.cambridge.org

Information on this title: www.cambridge.org/9781009605997

DOI: [10.1017/9781009605991](https://doi.org/10.1017/9781009605991)

© Xiao Lu 2026

This publication is in copyright. Subject to statutory exception and to the provisions
of relevant collective licensing agreements with the exception of the Creative
Commons version the link for which is provided below, no reproduction of any part
may take place without the written permission of Cambridge
University Press & Assessment.

An online version of this work is published at doi.org/10.1017/9781009605991
under a Creative Commons Open Access license CC-BY-NC-ND 4.0 which permits
re-use, distribution and reproduction in any medium for non-commercial purposes
providing appropriate credit to the original work is given. You may not distribute
derivative works without permission. To view a copy of this license, visit
<https://creativecommons.org/licenses/by-nc-nd/4.0>

When citing this work, please include a reference to the DOI [10.1017/9781009605991](https://doi.org/10.1017/9781009605991)

First published 2026

A catalogue record for this publication is available from the British Library

*A Cataloging-in-Publication data record for this Element is available from the Library
of Congress*

ISBN 978-1-009-60597-7 Hardback
ISBN 978-1-009-60602-8 Paperback
ISSN 2398-4023 (online)
ISSN 2514-3794 (print)

Cambridge University Press & Assessment has no responsibility for the persistence
or accuracy of URLs for external or third-party internet websites referred to in this
publication and does not guarantee that any content on such websites is, or will
remain, accurate or appropriate.

For EU product safety concerns, contact us at Calle de José Abascal, 56, 1º, 28003
Madrid, Spain, or email eugpsr@cambridge.org

Quantile Modeling for Political Research

Elements in Quantitative and Computational Methods for the Social Sciences

DOI: 10.1017/9781009605991
First published online: May 2026

Xiao Lu
Peking University

Author for correspondence: Xiao Lu, xiaolu@pku.edu.cn

Abstract: Quantile models are widely used across the natural and social sciences to analyze heterogeneous phenomena that conventional mean-based approaches often obscure. Yet, despite their growing importance in many disciplines, their adoption in political science has remained comparatively limited, in part because the field still lacks an accessible introduction tailored to its substantive questions and empirical practices. This Element addresses that gap by showing how quantile models can expand the methodological repertoire of political science and deepen our understanding of political phenomena. Combining methodological innovation with practical guidance, this Element introduces quantile models for both continuous and discrete response variables and illustrates their use with real-world political examples. All empirical applications are accompanied by publicly available data, code, and software, making the Element a useful resource for both teaching and research. This title is also available as Open Access on Cambridge Core.

Keywords: quantile regression, political science, political heterogeneity, hypothesis testing, computational social science

© Xiao Lu 2026

ISBNs: 9781009605977 (HB), 9781009606028 (PB), 9781009605991 (OC)
ISSNs: 2398-4023 (online), 2514-3794 (print)

Contents

1	Introduction	1
2	Quantile Models for Continuous Responses	14
3	Quantile Models for Binary Choices	33
4	Quantile Models for Multiple Choices	46
5	A Brief Overview of Other Quantile Methods	64
6	Conclusion	77
	References	80

1 Introduction

1.1 Motivation

Classical linear-regression models are typically used to estimate the average change in a response variable associated with a unit change in an explanatory variable. These mean-based models have dominated empirical political research for decades because of their parsimony, computational convenience, and relative ease of interpretation. Yet their standard inferential framework is often most straightforward under assumptions such as homoscedasticity and, in some settings, normally distributed errors. These assumptions may be problematic when the outcome distribution is skewed or when dispersion varies systematically across values of the covariates. More fundamentally, mean-based models focus attention on the conditional mean of the response variable given a set of covariates. As a result, heterogeneity away from that conditional mean is typically absorbed into the error term rather than treated as substantively meaningful variation in its own right.

In many political settings, however, differences across individuals, parties, governments, or institutions reflect genuine heterogeneity in preferences, incentives, constraints, or strategic environments. Such differences are often central to political analysis and cannot be adequately understood as mere random deviations from an average relationship. Consequently, an exclusive focus on averages may provide only a partial account of many substantive research questions in political science.

Indeed, many research questions are concerned less with average effects than with how political behavior varies across groups, time periods, institutional settings, or stages of strategic interaction (Braumoeller, 2006; De Marchi & Laver, 2023; Lu, 2020; Rosenberg, Knappe, & Braumoeller, 2017; Weschle, 2019). In such cases, reliance on average effects alone can produce incomplete and, at times, potentially misleading conclusions (Winship & Mare, 1992). Even when mean-based models allow for heteroscedasticity, they typically treat it as a problem for statistical inference to be corrected rather than as substantively meaningful variation to be examined in its own right. However, for many research questions, heterogeneity is precisely what matters. Units facing the same formal choice may behave differently because they are embedded in different contexts, constrained by different institutions, or guided by different preferences and incentives, including factors that are only imperfectly observed.

Quantile models, introduced in their modern form by Koenker and Bassett (1978), provide a powerful framework for analyzing heterogeneity in political data. Their intellectual roots reach further back: in 1760, Roger Boscovich

posed to Thomas Simpson a problem involving the minimization of the sum of absolute residuals, anticipating what would later be recognized as median regression (Stigler, 1984). Two centuries later, this line of thinking helped lay the foundation for modern quantile methods, which are now widely used across the natural and social sciences. While mean-based models focus on how covariates shift the conditional mean of a response variable, quantile models examine how relationships vary across different points of the conditional outcome distribution. This makes them especially useful for studying heterogeneous effects that may be obscured by average-based analysis. Rather than reducing variation away from the mean to residual noise, quantile models allow researchers to investigate how explanatory variables matter differently across the distribution of the response variable. They are also less sensitive to extreme values and offer a richer picture of the underlying data-generating process than models centered solely on average effects. Quantile models therefore broaden the range of political questions that can be addressed empirically, particularly when heterogeneity is itself of substantive interest.

These features make quantile models especially attractive for political scientists whose research questions are not well captured by conventional mean-based approaches. However, despite the aforementioned advantages, quantile models have been underutilized in political science until recently. Recognizing the utility of quantile models, scholars have increasingly applied them to examine political topics as diverse as geopolitical risks (Caldara & Iacoviello, 2022), political extremism (Bertelli & Richardson, 2008; Makowsky & Miller, 2014), coalition building (Häge, 2013), electoral change (Guinjoan & Rodon, 2021; Hill, Hopkins, & Huber, 2021), voter turnout (Kaniovski & Mueller, 2006; Schafer & Holbein, 2020), policy communication (Haan et al., 2022), political representation (Weschle, 2019), trade politics (Betz, 2017), foreign aid (Okada & Samreth, 2012), public budgeting (Breunig, 2011; Breunig & Jones, 2011; Breunig & Koski, 2020), interest group ratings (Brunell et al., 1999), executive constraints in parliaments (Abramson & Boix, 2019), political capital in developing countries (Appleton et al., 2009), political economy of natural disasters (Neumayer, Plümper, & Barthel, 2014), gender inequality in political participation (Arvate, Firpo, & Pieri, 2021), anchoring heuristic in public preference formation (Arceneaux & Nicholson, 2024), effects of wealth on political selection (Poulos, 2019), impact of democratization on corruption (Jetter, Agudelo, & Hassan, 2015), confirmation delay in the executive appointment process (Krause & Byers, 2022), grant retrenchments by US federal agencies (Krause & Zarit, 2022), economic consequences of terrorism on regional growth (Blomberg, Broussard, & Hess, 2011), distributional

inequity in public service provision (Cheng, Yang, & Deng, 2022), citizen responsiveness to government performance (Holbein, 2016), and effects of political institutions on private investment (Stasavage, 2002), among others.

Despite these applications, there are still several reasons that deter a wider application of quantile methods in the political science community. These mainly include (1) limited familiarity with quantile models due to the absence of a systematic introduction to the quantile approach in political science, (2) technical challenges in implementing quantile models and analyzing discrete choices in political settings, and (3) difficulties in presenting and interpreting results from quantile models due to multiple sets of coefficients for each quantile of interest.

This Element is designed to bridge the gap and provide political scientists with useful information about the most important aspects of quantile models. Throughout the Element, it becomes evident that the three primary obstacles in utilizing quantile models are straightforward to overcome. First, by explaining the basic concept of quantiles and quantile functions, I illustrate that the idea of quantiles is as intuitive as that of the mean, yet it offers greater flexibility in empirical analysis. Second, there is now an abundance of software packages and computational algorithms for implementing quantile methods in both continuous and discrete variable settings that remove the technical barriers to using quantile models. Applied political researchers can readily adopt existing quantile algorithms and software for their own research purposes. Third, although scholars unfamiliar with quantile regression might find it cumbersome to interpret results from quantile models – since each quantile requires its own interpretation and potentially its own graphical representation – this feature is what makes quantile models so useful. They provide a more complete picture of the relationship between variables by estimating effects at different points in the distribution of the response variable. Thus, the quantile approach offers significant advantages in terms of the ability to uncover nuanced insights across the distribution of the response variable. This benefit makes it a powerful tool for researchers interested in exploring the full complexity of their data.

Take vote choice as an example. Suppose we want to study how the ideological distance between voters and candidates influences voter decisions. A conventional mean-based model will summarize this relationship with a single average effect. However, it is entirely plausible that voters react differently to the ideological distance depending on their underlying propensity to vote for a certain candidate. In this case, unlike mean-based models, quantile models can capture this heterogeneity by showing how the effect of ideological distance

varies across different points of the conditional distribution of support, rather than restricting attention to the mean alone. Substantively, this has important implications for the strategic positioning of political candidates within the policy spectrum. If ideological distance has different effects among voters with lower and higher propensities to support, then candidates' strategic positioning in the policy space may influence core supporters and more persuadable voters in different ways. Subsequent sections will return to this and other political examples to illustrate how quantile effects can be interpreted and how quantile models can contribute to substantive political analysis.

In general, the *Element* is written as a combination of methodological innovation and practical guidance on quantile models. It introduces the basic properties of quantiles, illustrates quantile inference with continuous response variables, and then explores the potential of quantile methods in analyzing discrete choices. This *Element* also emphasizes that quantiles are as intuitive as the mean, yet they offer greater flexibility in modeling. Compared to existing books on quantile regression, this *Element* draws attention to discrete choice data in both binary and multiple choice forms, which play an especially important role in empirical research of political science. In particular, it introduces discrete quantile models for data with varying choice alternatives and for the estimation of multiple conditional quantiles. With real-world political examples, published software packages and hands-on computational codes, this *Element* demonstrates the utility of quantile models for a wide field of political research, and shows that estimating and interpreting quantile models can be as simple as estimating and interpreting the classical linear regression.¹

1.2 Aim and Organization

The aim of this *Element* is to encourage the practical use of the quantile approach in political science by introducing the concepts, functions, and applications of both the conventional and newly developed quantile models. It builds on the prior work of quantile regression but emphasizes more on the ability of quantile models in dealing with discrete choices and examining competing political theories. The *Element* is not intended to provide a comprehensive overview of all existing quantile methods, which is an impossible mission for a single book. While quantile methods have been extensively discussed in a variety of publications such as the *Handbook of Quantile Regression* edited by Koenker, Chernozhukov, He and Peng (Koenker et al., 2017), and

¹ The packages and data used in the *Element* are already publicly accessible, and the codes are kept in the public GitHub repository: https://github.com/xiao-lu-research/Quantile_Modeling_for_Political_Research.

books written by Koenker (2005), Hao and Naiman (2007), Davino, Furno, and Vistocco (2014), Furno and Vistocco (2018), and Cleophas and Zwinderman (2021), among others, these works are mainly geared toward statisticians and scholars outside the field of political science. On the contrary, this Element is tailored to the political science community, providing fundamental information on the properties and usage of quantile methods, along with instructional political examples and hands-on computational codes to demonstrate their substantive applications in the field. Because most political scientists have been exposed to mean-based models such as OLS, logit and probit models, this Element also compares their performance with quantile models to better illustrate the advantages of the latter in various political settings.

This Element is oriented mainly toward students and scholars in political science who are interested in utilizing quantile methods for their own learning and research purposes. For reproducibility and research purposes, the Element provides codes written in the R programming language, a popular and easy-to-learn open-source software project for data analysis and statistical computing (R Core Team, 2021). The advantage of using R is that it is free, publicly accessible, and allows readers to examine the source codes of the packages and functions, and consequently adjust and extend the functionalities as needed for their own research. This Element is not intended to provide an introduction to the R programming, as there are already a variety of excellent resources and tutorials available. The software is accessible via the CRAN website, which provides both source and binary versions for different operating systems and computing environments, including Windows, Linux, and Macintosh.²

The Element is organized as follows. The rest of this section will show the use of quantiles by introducing the basic concepts and properties of quantiles, quantile functions, and conditional quantiles. In Section 2, the traditional quantile regression applied for response variables in a continuous scale will be introduced. Section 3 expands the application of quantile models to cases with binary response variables. In this section, the estimation procedure of the binary quantile model will be introduced. Following the discussion of Section 3, a more generalized discrete quantile model (the conditional binary quantile model) for multiple choices will be presented in Section 4. Section 5 will provide a brief discussion on some more quantile methods, including those for estimating quantile treatment effects and survival data. This section will also introduce how to perform variable selection and model comparison in quantile settings. Besides the advantages of quantile models, the summary subsection of

² See <https://cran.r-project.org/>.

the section will discuss practical issues of the quantile approach. The final section concludes by highlighting briefly why and when quantile models should be used.

1.3 Basic Concept of Quantiles

Generalizing notions such as median, quartile, decile, and percentile, quantiles are a set of values that divide a distribution into ordered parts. The τ th quantile of a given set usually denotes the smallest value below which at least a proportion τ of the observations fall. One of the most well-known quantiles is the median (or the 0.5th quantile), which separates the lower half from the higher half of a distribution. Other widely used quantiles include the first and third quartile, which are also known as the 0.25th and 0.75th quantile, or the 25th and 75th percentile. They are commonly used to calculate the interquartile range as a measure of data dispersion.

As quantiles are only sensitive to the ordering of the data but not to extreme values, they represent a robust alternative to the mean when evaluated at the center of a distribution. They can also offer a more informative summary of dispersion when the data are skewed or otherwise depart from normality. To illustrate these basic properties, consider for example in Figure 1 a set of values [1, 2, 3, 7, 8, 9, 10, 11, 12, 13, 100] arranged in ascending order. This set is clearly not normally distributed, but right-skewed in the sense that most observations are concentrated at lower values, with a long right tail. The 0.25th quantile, or equivalently the first quartile, of the set is the value below which there are at least 25% of cases, and in this example falls between 3 and 7 under common sample definitions. In the same vein, we can calculate the 0.75th quantile (third quartile), which is located between 11 and 12. Due to the existence of an extreme value (100) in this example, it is also apparent that compared to the mean 16, the median 9 is a more informative summary of the central location. These values can then be used to quantify the dispersion of the sample. Compared to the dispersion calculated by the sample standard deviation (approximately 28), which is more than twice the difference between the smallest and the second largest value of the sample, the interquartile range measured by the difference between the 0.25th and 0.75th quantile, as shown by the box plot in Figure 1, represents a more meaningful summary of the dispersion of the sample. Moreover, we can calculate another robust alternative to the mean-based standard deviation – the median absolute deviation (MAD), which is defined as the median of the absolute deviations from the median of the sample $\mathbf{x} = \{x_1, x_2, \dots, x_n\}$:

$$\text{MAD} = \text{Med} [|x_i - \text{Med}(\mathbf{x})|] . \quad (1.1)$$




Figure 1 Sample quantiles and box-plot

The MAD of the previous sample is 3, representing a more representative summary statistic for variability of the sample than the standard deviation.

As the figure also illustrates, the standard box-plots use quantiles to summarize the central tendency, dispersion, and skewness of a dataset, while also helping identify outliers. More generally, denote the value at τ th quantile as q_τ . For any $\tau < 0.5$, the inter-quantile range between the lower (τ) and the upper ($1 - \tau$) quantile can be computed as follows:

$$\Delta q = q_{1-\tau} - q_\tau, \quad (1.2)$$

which denotes the dispersion of the middle $(1 - 2\tau) \times 100\%$ of the sample.

Because of their robustness and their ability to characterize different parts of a sample or distribution, quantiles have become common statistics in descriptive summaries, quantification of uncertainty, and hypothesis testing in empirical research. A familiar example is the quantile-to-quantile (QQ) plot, which is commonly used to assess whether an empirical distribution is approximately normal. By comparing the quantiles of the dataset with the theoretical quantiles of a normal distribution, the QQ plot provides a simple visual diagnostic of deviations from normality.

1.4 Cumulative Distribution Function and Quantile Function

The cumulative distribution is closely related to the concept of quantiles. To understand quantiles more formally, it is thus helpful to clarify the relation between the cumulative distribution function and the quantile function. For any random variable Y , the cumulative distribution function evaluated at y is defined as the probability that Y is less than or equal to y :

$$F_Y(y) = \Pr(Y \leq y). \quad (1.3)$$

Statistically, scholars may be interested in knowing the location of a specific value within a distribution besides the mean. Consider the cumulative distribution function of the standard normal distribution as shown in Figure 2a.




Figure 2 Cumulative distribution function (a) and quantile function (b)

We might wish to know, for example, what proportion of the distribution lies below the values -0.53 and 0.84 . To do so, we trace each value on the horizontal axis up to the CDF and then across to the vertical axis. The resulting values are approximately 0.3 and 0.8 , respectively (Figure 2a). In other words, -0.53 corresponds to the 0.3 th quantile and 0.84 to the 0.8 th quantile of the standard normal distribution. The quantile function, as shown in Figure 2b, reverses this mapping by returning the value associated with a given cumulative probability.

Formally, the τ th quantile function of a random variable Y , denoted $Q_Y(\tau)$, is defined as the generalized inverse of its cumulative distribution function:

$$Q_Y(\tau) := F_Y(\tau)^{-1} = \inf\{y : F_Y(y) \geq \tau\}. \quad (1.4)$$

The quantile function is useful because it allows us to recover values associated with specified probability levels. This is central to many tasks in statistical analysis, including the construction of critical values, confidence intervals, prediction intervals, and other forms of uncertainty assessment.

Although symmetrical distributions such as a normal distribution can often be summarized reasonably well by its central tendency and dispersion, that is, its mean and variance, the mean-based statistics become less informative when the distribution is asymmetric. Consider for instance a right-skewed distribution whose cumulative distribution function is plotted in Figure 3.³ Approximating the distribution using a normal distribution with the corresponding mean and standard deviation will misrepresent the distribution around the median and in the tails. On the contrary, quantile approximation uses piecewise information from the random sample, which, as shown by the dashed lines in the figure, yields a much better fit.

³ Here, we illustrate an Asymmetric Laplace Distribution with location parameter $\mu = 0$, scale parameter $\lambda = 1$, and skewness parameter $\kappa = 50$.




Figure 3 Approximation of a right-skewed distribution (asymmetric Laplace distribution with location parameter $\mu = 0$, scale parameter $\lambda = 1$, and skewness parameter $\kappa = 50$)




Figure 4 Summary of the distribution of ethnic vote shares by mean (a) and quantile (b)

The substantive value of quantile-based summaries becomes especially clear when political outcomes are distributed unevenly across units. Take the distribution of ethnic voting in Bosnia as an example. Figure 4 illustrates the distribution using the violin plot. Based on data collected by Hadzic, Carlson, and Tavits (2020), the average municipal-level vote share of ethnic parties in national legislative elections is 80.99% (see Figure 4(a)). Compared to the median share of 84.89% (see Figure 4(b)), the mean value is a reduction of 4 percentage points. This indicates that the whole sample is distributed asymmetrically, with observations concentrated at relatively high vote shares and a longer tail toward lower values. In addition to the skewness, the spread

of the distribution is wide, with the smallest share of 19.33% and the largest share of 99.49%.

Moreover, as the left panel of the figure shows, if we use the 95% interval based on the normality assumption to quantify the spread of the variable, the upper limit of the 95% interval will exceed even the highest value of the sample, while the lower limit will underestimate the lowest values. Furthermore, because the quantification of the spread is symmetric around the mean, the mean-based approach is unable to reveal the skewness of the variable. In contrast, by examining the 0.95th and 0.05th quantile as illustrated in the right panel of Figure 4, we can find a sharp distinction between the upper and lower halves of the distribution: While there is only a 13.15% difference between the 0.95th quantile and the median, the difference between the median and the 0.05th quantile is 30.83%, more than twice the difference in the upper half. A focus on averages alone would therefore obscure substantively important variation across municipalities in the distribution of ethnic voting.

1.5 Conditional Quantiles

The illustrations in Section 1.4 show that quantiles provide added value to the mean by offering a more complete summary of a distribution. However, in political research, the central interest rarely lies in the distribution of a single variable alone. Instead, we want to investigate the effect of variables on the outcome of interest, that is, how changes in explanatory variables are associated with changes in the outcome. To reveal the conditional relationship between variables, we need to compute conditional quantiles. Conditional quantiles are “conditional” in the sense that the estimates depend on the covariates on the right-hand side of the estimation equation, and thereby they reveal local effects of a variable on the response across the whole distribution.

To show the benefits of using conditional quantiles, recall the standard linear model:

$$y_i = \mathbf{x}_i' \boldsymbol{\beta} + \epsilon_i. \quad (1.5)$$

If we assume that the conditional expectation of the error term equals zero (i.e., $E(\epsilon_i | \mathbf{x}_i) = 0$), we obtain the conditional mean model. Subsequently, we can reformulate the equation with the response variable y on the left-hand side and a vector of explanatory variables $\mathbf{x}$ on the right-hand side (Lewis-Beck & Lewis-Beck, 2015):

$$E(y_i | \mathbf{x}_i) = \mathbf{x}_i' \boldsymbol{\beta}. \quad (1.6)$$

This formulation summarizes the relationship between $\mathbf{x}_i$ and y_i at the conditional mean of the outcome. As a consequence, the covariate effect is

represented by a single coefficient vector β , rather than being allowed to vary across different parts of the conditional distribution. The coefficients can then be estimated by solving:⁴

$$\hat{\beta} = \underset{\beta}{\operatorname{argmin}} \sum_{i=1}^N (y_i - \mathbf{x}_i' \beta)^2, \quad (1.7)$$

the familiar least-squares problem. While highly useful, the conditional mean neglects important information about potentially heterogeneous effects of $\mathbf{x}$ on y .

On the contrary, if we introduce an alternative formula

$$Q_{\tau}(y_i | \mathbf{x}_i) = \mathbf{x}_i' \beta_{\tau}. \quad (1.8)$$

by replacing the expectation $E(\cdot)$ with conditional quantiles $Q_{\tau}(\cdot)$, we are in a more flexible modeling situation. In this formula, each coefficient vector β_{τ} represents conditional effects at a specific quantile location τ of the response variable (for a formal proof, see e.g. Ding (2025, 294–297)). For $\tau = 0.5$, this becomes the conditional median model, which can be estimated by minimizing the sum of the absolute deviations:

$$\hat{\beta}_{0.5} = \underset{\beta}{\operatorname{argmin}} \sum_{i=1}^N |y_i - \mathbf{x}_i' \beta|. \quad (1.9)$$

More generally, quantile regression estimates β_{τ} by minimizing the asymmetric absolute loss:

$$\hat{\beta}_{\tau} = \underset{\beta}{\operatorname{argmin}} \sum_{i=1}^N \rho_{\tau}(y_i - \mathbf{x}_i' \beta), \quad (1.10)$$

where $\rho_{\tau}(x) = x[\tau - I(x < 0)]$ (for details of derivation, see Section 2.2). This objective function, often called the check function, gives asymmetric weights to positive and negative residuals and thereby identifies the conditional quantile.

Formally, the conditional quantile value of y_i given $\mathbf{x}$ can be written as

$$q_{\tau} | \mathbf{x}_i = \inf\{y_i : F(y_i | \mathbf{x}_i) \geq \tau\}, \quad (1.11)$$

where $F(\cdot)$ is the conditional cumulative distribution function. For each quantile level τ , we are able to get a quantile-specific effect β_{τ} that may differ from the average effect.

To interpret β_{τ} , we can regard it as the marginal change in the τ th quantile of the response variable due to a marginal change of the explanatory variable(s).

⁴ While the variance of the conditional means can capture the deviation of the estimated effect by heterogeneous subgroups, it still remains unclear about the direction and magnitude of varying subgroup effects.

In causal applications, and under additional identification assumptions, related quantile-based parameters can also be interpreted in terms of treatment effects at different points of the outcome distribution (Frandsen, Frölich, & Melly, 2010; Frölich & Melly, 2012), the details of which will be discussed in the subsection on quantile treatment effects in Section 5.

To illustrate the contrast between conditional means and conditional quantiles, return to the aforementioned example of ethnic voting. Suppose we are interested in the effect of wartime violence on ethnic voting, measured by wartime casualties and ethnic vote share, respectively (Hadzic, Carlson, & Tavits, 2020). In a simple bivariate specification, the conditional mean can be calculated using the OLS regressing (logged) casualty against ethnic vote share. As illustrated by the solid line and the associated 95% confidence interval in Figure 5, the estimated mean relationship is not statistically distinguishable from zero. Once we examine the relationship across quantiles, however, a more differentiated pattern emerges. As shown by the fitted lines at the 0.05th, 0.5th, and 0.95th conditional quantiles in the figure, the positive relation between wartime violence and ethnic voting is strongest in lower quantiles of ethnic vote distribution and weakens as one moves toward the upper quantiles.




Figure 5 Effects of wartime violence on ethnic voting

This points to substantively meaningful heterogeneity across municipalities, possibly reflecting differences in prior levels or traditions of ethnic voting. Examining conditional quantiles alongside the conditional mean thus provides a fuller and more nuanced account of the relationship between wartime violence and ethnic voting.

Note that for illustrative purposes, the depicted effects at this stage are solely based on the bivariate analysis, which consequently neglects potential confounding variables. In the [following section](#), we will return to this example with a more complete analysis controlling for potential confounders.

1.6 Summary

This section has introduced the basic features of quantiles, the role of quantile functions, and the logic of conditional quantiles. According to the above illustrations, [Table 1](#) summarizes the main differences between the mean-based and the quantile approach.

The quantile approach can be used to achieve the same goal as the mean-based approach, that is, to represent the central tendency of a sample or distribution. Compared to the mean, the median is more robust to extreme values. While the mean-based approach is often most straightforward under assumptions such as symmetry, homoscedasticity, or normally distributed errors, the quantile approach allows for more flexible distributions and relaxes the conditional-zero-mean assumption of the error term. When data contain heterogeneous units or the distribution of interest is skewed and asymmetric, quantiles provide additional information that would be overlooked when using the mean-based approach.

Table 1 Comparison between mean-based and quantile approach

	Mean-based approach $E(y_i x_i) = x_i'\beta$	Quantile approach $Q_\tau(y_i x_i) = x_i'\beta_\tau$
Central tendency	Mean	Median
Data distribution	Normality assumption	Without normality assumption
Error term	$E(\epsilon x) = 0$	$Q_\tau(\epsilon_\tau x) = 0$
Spread	Standard deviation	Median absolute deviation
Estimated effect	Single average effect	Multiple local effects across quantiles
Estimation	Least squares	(Weighted) Least absolute deviation

Mean-based approaches often summarize dispersion with the standard deviation and typically rely on variance-based measures for statistical inference. Without appropriate adjustments, this will lead to biased estimates of standard errors in the presence of heteroscedasticity, that is, when the variance of the error term is not constant. The quantile approach overcomes this problem by estimating local effects at different quantiles. The power of quantiles is not only about the complete description of data or distributions, but also the ability to investigate conditional relations between variables. This is particularly useful when we want to know how changes of the explanatory variables affect the outcome across its entire distribution. Therefore, in addition to robustness checks, quantile models allow researchers to examine distributional effects among heterogeneous political units, which will provide a more complete picture of the relationship between variables of interest.

This does not mean that quantile methods replace mean-based analysis in all applications. Rather, they complement it by expanding the range of questions that can be asked and answered empirically. Building on these foundations, the sections that follow extend the discussion to multivariate settings and show how quantile methods can be applied to a variety of statistical and substantive problems in political science.

2 Quantile Models for Continuous Responses

2.1 Introduction

This section introduces the classical quantile model, which focuses on continuous response variables. In contrast to the univariate setting discussed in the [previous section](#), it demonstrates how to estimate conditional quantiles for the continuous response variable in multivariate settings. In continuous scales, the response variable is represented by values on the real number line, providing rich information commonly seen in empirical political research. Variables of this kind, such as vote shares, party size, and level of turnout, carry substantive meanings through differences in magnitude. Modeling continuous response variables usually eliminates the need for additional data transformations that are common for evaluating discrete response variables discussed in the following chapters. Consequently, the coefficients obtained from quantile models directly reflect how the response variable changes when the explanatory variables shift by one unit at the specified quantile location.

Formally, adopting linear parameterization, quantile regression models the data with the continuous response variable y_i at τ th quantile as

$$Q_\tau(y_i|x_i) = x_i'\beta_\tau, \quad (2.1)$$

where $\mathbf{x}$ corresponds to the variables of the data, and $\boldsymbol{\beta}_\tau$ is a vector of coefficients associated with the variables at quantile τ . The core task of quantile regression is to estimate values of $\boldsymbol{\beta}_\tau$ given $\mathbf{y}$ and $\mathbf{x}$, the details of which will be discussed in the following subsection.

2.2 Estimation and Inference in Quantile Regression

In contrast to the conditional mean-based approach that minimizes a squared loss function, the conditional τ th quantile as a solution to the above equation is obtained by minimizing an asymmetric loss function $\mathcal{L}$, where each component is weighted either by τ or $1 - \tau$ (Yu & Moyeed, 2001):

$$\mathcal{L} = \tau \int_{y > q} |y - q| dF_Y(y) + (1 - \tau) \int_{y < q} |y - q| dF_Y(y). \quad (2.2)$$

In this formula, $F(\cdot)$ is a cumulative distribution function, and q represents the value of the τ th quantile as calculated by Equation (1.4). According to Equation (2.2), the quantile loss function uses all the sample information, instead of splitting the sample, to derive the quantile values. Because the linear loss function depends not on certain covariate values but on covariate orderings, the quantile estimators are more robust than the least squares estimators in the presence of extreme values.

By solving the first order condition, we have the τ th sample objective function evaluated at q as

$$\frac{1}{N} \sum_{i=1}^N \rho_\tau(y_i - q), \quad (2.3)$$

where

$$\rho_\tau(x) = x [\tau - I(x < 0)] = [(1 - \tau)I(x < 0) + \tau I(x \geq 0)] |x| \quad (2.4)$$

is known as the check function and $I(\cdot)$ denotes the indicator function. As illustrated in Figure 6, the check function takes different shapes when different quantiles are specified. The function is symmetric when evaluated at the median (0.5th quantile shown by the solid line).

Given a linear model $y_i = \mathbf{x}_i' \boldsymbol{\beta} + \varepsilon_i$, the τ th quantile regression estimator minimizes the above equation by replacing q with $\mathbf{x}_i' \boldsymbol{\beta}_\tau$. In the model, we commonly assume $Q(\varepsilon_i | \mathbf{x}_i, \tau) = 0$. If $\tau = 0.5$, the model reduces to a median regression. The goal is to estimate the values of $\boldsymbol{\beta}_\tau$ for each τ by solving the following minimization problem:

$$\hat{\boldsymbol{\beta}}_\tau = \underset{\boldsymbol{\beta}}{\operatorname{argmin}} \sum_{i=1}^N \rho_\tau(y_i - \mathbf{x}_i' \boldsymbol{\beta}). \quad (2.5)$$




Figure 6 Check functions at different quantiles

The common solution to this minimization problem is based on the technique of linear programming, which iterates through points and lines on a polyhedral surface to find the optimal estimates. The standard errors and confidence intervals are then calculated based on the asymptotic normality assumption or by repeated sampling using bootstrap methods (Davino, Furno, & Vistocco, 2014; Koenker, 2005). It can be shown that under certain regularity conditions, the estimated quantile coefficients $\hat{\beta}_\tau$ are asymptotically normal:

$$\sqrt{n}(\hat{\beta}_\tau - \beta_\tau) \xrightarrow{L} N(0, \Lambda_\tau), \quad (2.6)$$

where $\Lambda_\tau = \tau(1 - \tau)(E[f_{\epsilon_\tau}(0|\mathbf{x}_i)\mathbf{x}_i\mathbf{x}_i'])^{-1}E[\mathbf{x}_i\mathbf{x}_i'](E[f_{\epsilon_\tau}(0|\mathbf{x}_i)\mathbf{x}_i\mathbf{x}_i'])^{-1}$ and $f_{\epsilon_\tau}(0|\mathbf{x}_i)$ is the conditional density of the error term evaluated at zero (Buchinsky, 1998).

Alternatively, recent developments in Bayesian methods allow researchers to estimate quantile regression using the Markov chain Monte Carlo (MCMC) sampling procedure (Yu & Moyeed, 2001). This is done by imposing appropriate (or uninformative) priors on the parameters to be estimated, and then combining the prior with the likelihood function to form the posterior distribution, which can be subsequently sampled using MCMC algorithms. Compared to the frequentist approach which relies on asymptotic properties, the advantage of the Bayesian estimation is that it provides a natural quantification of estimation uncertainty from random draws of the posterior distribution.

As proved by Koenker and Bassett (1978, 39) (Theorem 3.2 of the paper), there are a number of desirable equivariance properties of the estimated coefficients $\hat{\beta}$ that can be statistically interesting to applied researchers:

$$\hat{\beta}(\tau, \lambda y, X) = \lambda \hat{\beta}(\tau, y, X), \lambda \in [0, \infty), \quad (2.7)$$

$$\hat{\beta}(1 - \tau, \lambda y, X) = \lambda \hat{\beta}(\tau, y, X), \lambda \in (-\infty, 0], \quad (2.8)$$

$$\hat{\beta}(\tau, y + X\gamma, X) = \hat{\beta}(\tau, y, X) + \gamma, \gamma \in \mathbb{R}^K, \quad (2.9)$$

$$\hat{\beta}(\tau, y, XA) = A^{-1} \hat{\beta}(\tau, y, X), A_{K \times K} \text{ nonsingular}, \quad (2.10)$$

where the first two equations suggest that the coefficients are equivariant under scalar transformations of the response variable. This property is particularly useful for analyzing discrete response variables. The last two equations state that the coefficients are equivariant to a location shift of the response variable and reparameterization of the design matrix. Thus, it provides more flexibility for regression analysis.

To evaluate the goodness of fit of quantile regression, Koenker and Machado (1999) also developed an analog of the R -squared measure:

$$R_\tau = 1 - \hat{V}_\tau / \bar{V}_\tau, \quad (2.11)$$

where $\hat{V}_\tau = \sum_1^N \rho_\tau(y_i - \hat{y}_i)$ and $\bar{V}_\tau = \sum_1^N \rho_\tau(y_i - \bar{y}_i)$. Here, $\hat{y}_i$ and $\bar{y}_i$ represent respectively the predicted response value and the sample quantile of the response variable without controlling for covariates.

For applied researchers, fortunately, there are statistical software packages that can perform estimation tasks for us. So here I will not delve into the specifics of the various types of estimators and their solution concepts. To demonstrate the utility of quantile estimation tools for analyzing continuous response variables, we can utilize two open-source software packages written in the R language. The first is the `quantreg` package developed by Roger Koenker and continuously enhanced by numerous statisticians (Koenker, 2025). It offers a diverse set of functions for estimating conditional quantiles using a frequentist approach. The second package, `bayesQR`, was created by Dries Benoit, Rahim Al-Hamzawi, Keming Yu, and Dirk Van den Poel (Benoit & Van den Poel, 2017), primarily based on the methods proposed in Benoit and Van den Poel (2012) and Benoit, Alhamzawi, and Yu (2013). This package is used for Bayesian quantile regression estimation. To install and load these packages, we can execute the following codes, which download the package sources from the Comprehensive R Archive Network (CRAN) to the local drive and load the packages into the current working environment.

```
1 # Install two packages from CRAN repository
2 install.packages(c("quantreg", "bayesQR"))
3 # Load the packages
4 library(quantreg)
5 library(bayesQR)
```

In what follows, the quantile estimation will be illustrated using two real-world political examples of ethnic voting and political representation. For the estimation of the data, the analysis will mainly exploit the `rq()` function from the `quantreg` package (details on the `quantreg` package are preserved for the [next section](#)). Note that estimating the data of the two examples using the Bayesian approach without informative priors yields similar results.

2.3 Example of Ethnic Voting

To illustrate the quantile model, consider again the ethnic voting example studied by Hadzic, Carlson, and Tavits (2020). The authors argue that community-level experience with wartime violence increases the salience of ethnic identities, making it important for postwar vote choice. Employing a difference-in-differences design on a panel dataset from pre- and postwar elections of Bosnia, Hadzic, Carlson, and Tavits (2020) find strong support for a positive effect of wartime violence on postwar ethnic voting. Yet the theoretical logic of the argument also suggests that this effect may not be uniform across communities. Quantile regression provides a natural way to examine this possibility by asking whether the relationship between wartime violence and postwar ethnic voting differs across the lower and upper parts of the conditional outcome distribution. In this example, different conditional quantiles represent the varying proportions of postwar ethnic voting in different communities. For example, a conditional quantile of 0.1 would roughly represent the bottom 10% communities (about 10 communities in the sample) where the proportion of ethnic voting is lower than the rest 90% of the communities in the sample. Because communities with low proportions of ethnic voting in postwar elections are mostly those communities with low prewar ethnic voting, the quantile approach also allows us to examine the variation of ethnic voting via the lens of diverse cultures of cooperation and inter-ethnic relations in different communities.

In the literature on civil conflict, there are at least two competing theories explaining the impact of wartime violence on the politicization of ethnicity. One theory posits that fears of violence and prewar norms of cooperation will reduce politicization of ethnicity after the war (Bellows & Miguel, 2009; Blattman, 2009; Gilligan, Pasquale, & Samii, 2014; Voors et al., 2012; Wood, 2003). This could be attributed to the possibility of more altruistic individuals remaining disproportionately in war-plagued communities, or to the need of the remaining populace to collectively confront threats and psychological distress (Gilligan, Pasquale, & Samii, 2014). Conversely, another theory emphasizes the heightened salience of ethnicity and the role of war-enhanced group identification in promoting postwar ethnic voting (Hadzic, Carlson, & Tavits, 2020;

Wilkinson, 2006).⁵ According to this theory, ethnic violence during wartime amplifies rather than diminishes inter-ethnic distrust, rendering ethnic parties a more appealing avenue for post-war representation (Hadzic, Carlson, & Tavits, 2020). However, the conflicting theories and their empirical evidence also indicate possible diversity among various subgroups of a population that may react differently to ethnic violence. For example, based on the level of prewar social cohesion, the influence of exposure to ethnic violence during wartime on ethnic voting may fluctuate across different communities. It is plausible to expect that in regions where social cohesion is high and thus ethnic polarization is low, the experience of wartime violence is likely to reduce ethnic voting after the war. However, in regions where there is relatively high ethnic polarization, we may expect a positive effect of wartime violence on ethnic voting. If such heterogeneity exists, revisiting the data with quantile regression could potentially unveil varying subgroup effects that have significant implications but have been largely disregarded in the literature.

The data collected by Hadzic, Carlson, and Tavits (2020) record election results in Bosnia both before and after the Bosnian Civil War fought between 1992 and 1995, which are publicly available.⁶ After downloading the data, we can load and explore the detailed features of the variables with the following codes.

```

1 # Load the ethnic voting data
2 data <- read.csv(file="MainData.csv")
3 # View the names of all variables
4 names(data)
5 # Keep selected variables for illustration
6 data_explore <- subset(data, select = c(
7   "Municipality", "Year", "Ethnic_Vote_Share",
8   "Casualty", "Log_Casualty"))
# Display the first five observations
head(data_explore, 5)

```

As shown in Table 2, the response variable `Ethnic_Vote_Share` measures the municipal-level percentage of votes cast for ethnic parties and varies over time. More specifically, the data come from 107 Bosnian communities,

⁵ Some other research also indicates that the effect of violence is contingent on the distinction between in- and out-group members (De Luca & Verpoorten, 2015; Rohner, Thoenig, & Zilibotti, 2013a, 2013b). For instance, Grossman, Manekin, & Miodowni (2015) suggest that only in-group members exhibit high social cohesion, trust, and altruism after violence exposure.

⁶ The data is publicly accessible through the Harvard Dataverse using the link <https://dx.doi.org/doi:10.7910/DVN/EUAIFA>.

Table 2 First five observations from subset of ethnic voting data

	Municipality	Year	Ethnic_Vote_Share	Casualty	Log_Casualty
1	Velika Kladusa	2006	21.15	2.05	0.72
2	Velika Kladusa	2010	19.33	2.05	0.72
3	Velika Kladusa	2014	33.37	2.05	0.72
4	Cazin	2006	64.84	1.47	0.38
5	Cazin	2010	61.95	1.47	0.38
:	:	:	:	:	:

including Velika Kladusa and Cazin in years 1990, 2006, 2010, and 2014. The casualty measure records the intensity of ethnic violence and is taken from the Bosnian Book of the Dead, which records the percentage of the total number of confirmed dead and missing against the prewar municipal population (Hadzic, Carlson, & Tavits, 2020, 350). The variable `Log_Casualty` is the log-transformed value of `Casualty`. Using these measures, we are able to first explore the primary relation between (logged) casualty and ethnic vote share using scatter plots.

```
1 # Scatter plot of Casualty and Ethnic_Vote_Share
2 plot(x = data_explore$Casualty, y = data_explore$
      Ethnic_Vote_Share, pch = 18, col = "gray",
      xlab = "Casualty", ylab = "Ethnic Vote
      Share")
3 # Scatter plot of Log_Casualty and Ethnic_Vote_
  Share
4 plot(x = data_explore$Log_Casualty, y = data_
      explore$Ethnic_Vote_Share, pch = 18, col =
      "gray", xlab = "Casualty (Logged)", ylab =
      "Ethnic Vote Share")
```

The scatter plots in Figure 7 reveal that the original measure of casualty is right-skewed (left panel), but is close to a normal distribution in the logarithmic scale (right panel). Therefore, as with Hadzic, Carlson, and Tavits (2020), the logged version of the casualty measure is adopted for the regression analysis. The `quantreg` package is used to estimate the conditional quantiles, with the parameter `tau` specifying the quantiles to be estimated. For the values of `tau`, a range of quantiles from 0.1 to 0.9 are specified. Here, lower quantiles may be interpreted as pertaining to communities where the proportion of ethnic voting is relatively low, and upper quantiles represent those communities where votes are more divided along ethnic lines. Given control variables, conditional quantiles can also be interpreted as capturing different positions in the




Figure 7 Scatter plots of correlation between casualty(a)/logged casualty (b) and ethnic vote share

distribution of ethnic vote share across communities, which may reflect varying degrees of ethnic political alignment. This allows us to ask whether the association between wartime violence and ethnic voting differs systematically across communities located at different points of the ethnic-voting distribution.

For illustrating the quantile estimation of the effects of wartime violence on ethnic voting, let us first consider a simple model of the relation between the two variables. A more comprehensive model in the subsequent analysis will also take into account the spatial and temporal variation. Here, the `rq` function from the `quantreg` package is used (Koenker, 2025), and the computational codes are shown in the following. The results can be visualized and printed using the functions `plot` and `summary`.

```
1 # Estimate the quantile model at quantiles from
  0.1 to 0.9
2 mod1 <- rq(Ethnic_Vote_Share ~ Log_Casualty, data
  = data, tau = seq(0.1,0.9,0.1))
3 # Plot point estimates
4 plot(mod1)
5 # Summarize the estimation results with
  confidence intervals
6 summary(mod1)
```

Figure 8 shows the point estimates of the effects for the intercept and casualty at different quantiles, where the dashed horizontal lines represent the OLS estimates. In both panels of the figure, the x-axis represents different quantile levels, and the y-axis shows the estimated effects of a variable at different quantiles, with the dotted line indicating how the effects change with increasing quantile levels. Compared to a single average effect estimated by the OLS, the U-shaped quantile estimation curve of casualty in the lower panel demonstrates that the effects of casualty on ethnic voting may differ across subgroups. In the lower quantiles less than 0.6 (except for 0.1), the effects are negative, while in




Figure 8 Point estimates of quantile effects for ethnic voting example

Note: In both panels, the x-axis represents different quantile levels, and the y-axis shows the estimated effects of a variable at different quantiles. Black dotted lines represent quantile estimates and horizontal dashed lines represent OLS estimates. The upper panel shows the quantile effects of the intercept, while the lower panel shows the quantile effects of the logged casualty.

the upper quantiles higher than 0.6, the effects become positive. These results reveal significant differences in the effects of casualty on ethnic voting between communities with diverse levels of ethnic voting in wartime. Because quantiles are measured on shares of ethnic votes (and thus ethnic polarization) across communities, it indicates that exposure to violence may affect ethnic voting differently depending on the ethnic polarization in each community. In particular, in lower quantiles where ethnic polarization is relatively low, the effects of wartime violence (*Log_Casualty*) on ethnic voting become negative, which suggests that in communities with low ethnic polarization, exposure to ethnic violence is likely to increase inter-ethnic trust and postwar social cohesion (and thus reduces the proportion of ethnic voting). This is the case for Bosnian communities such as Centar Sarajevo, Novo Sarajevo, Srebrenik, Tuzla, and Velika Kladusa, where the average proportion of ethnic voting is lower than the 0.1th quantile of the whole sample (i.e., 65% ethnic vote share). However, as demonstrated by the positive effects in the upper quantiles, the opposite is true in other communities where ethnic polarization is high. Overall, the results show that in lower quantile regions where the proportion of ethnic voting is relatively low, the OLS is likely to overestimate the effect of casualty, while in upper quantiles it is more likely to underestimate the effect.

The simple bivariate model is useful for illustration, but it leaves open the possibility that the estimated pattern is confounded by differences across years and municipalities. To better account for potential confounding factors, we now examine a more comprehensive model that takes into account the spatial and temporal variation. More specifically, we follow the model specification and the difference-in-differences (DID) design of Hadzic, Carlson, and Tavits (2020), with dummy controls of years before and after the civil war. The analysis also includes the fixed effects of regions. The following codes show the details of the quantile estimation procedure for the ethnic voting data.

```

1  # DID design: Generate dummy matrix
   distinguishing between years before and after
   war
2  dummy.matrix <- as.matrix(cbind(as.numeric(dat_
   violence$Year == 2006), as.numeric(dat_
   violence$Year==2010), as.numeric(dat_violence$
   Year==2014)))
3  # Quantile estimation with municipality-fixed
   effects
4  mod2 <- rq(Ethnic_Vote_Share ~ Log_Casualty:dummy
   .matrix+dummy.matrix+Municipality-1, data =
   data, tau = seq(0.1,0.9,0.1))
5  # Summarize the results
6  mod2_est <- summary(mod2)
7  # Extract coefficient estimates of the main iv.
   while omitting others
8  extract_coef <- function(x) return(x$coefficients
   [111:113,])
9  # Put coefficients into a single data frame
10 coef_df <- as.data.frame(do.call("rbind",lapply(
   mod2_est, extract_coef)))
11 coef_df$Year <- rep(c(2006,2010,2014),9)
12 coef_df$quantile <- rep(seq(0.1,0.9,0.1),each =
   3)
13 names(coef_df)[2:3] <- c("lower","upper")
14 # Visualize the coefficients
15 library(ggplot2)
16 ggplot(coef_df,aes(x = quantile, y = coefficients
   ))+
17 geom_line()+
18 geom_ribbon(aes(ymin = lower, ymax = upper),alpha
   = 0.2)+
19 facet_wrap(~Year)+
20 geom_hline(yintercept = 0,linetype = "dashed")+
21 xlab("Quantiles")+
22 ylab("Effect of Wartime Violence on Ethnic Vote
   Share")+
23 theme_bw()

```




Figure 9 Estimated effects of wartime violence on ethnic vote share over postwar years and across quantiles

Note: Areas surrounding the lines represent the 95% confidence interval.

The results of the coefficient estimates of the OLS and quantile regression are displayed in Figure 9, which shows the results in 2006 (a – left panel), 2010 (b – middle panel), and 2014 (c – right panel) in comparison to the baseline year 1990 in the prewar period. The OLS estimation follows Model 1 in table 1 of Hadzic, Carlson, and Tavits (2020, 353) with clustered (on municipality and year) robust standard errors. Compared to the average positive estimates by the OLS, all three panels of the figure show that while the positive effects of wartime violence on ethnic voting are evident in the upper quantiles where ethnic voting is more common, the effects are insignificant in the lower quantiles (quantiles lower than 0.5). Compared to the results in 2006 and 2010, the reduced effect of wartime violence on ethnic voting in the middle and upper quantiles in 2014 also indicates that the impact of wartime memory of ethnic violence seems to be decreasing at the community level.

Given that communities with low proportions of ethnic voting in postwar elections are mostly those communities with low prewar ethnic voting, it is plausible to associate the level of ethnic voting with the culture of cooperation and inter-ethnic relations in different communities. For example, as shown in Table 3, for communities such as Centar Sarajevo, Novo Sarajevo, Srebrenik, and Tuzla, where wartime violence has no discernible effects, they are all below the 0.1th quantile (i.e., around 65%) measured by the proportion of

Table 3 Quantile estimates and fixed-effects of four communities at 0.1th quantile

Variable	Coefficients	Lower bound	Upper bound
Ethnic Vote Share (2006)	2.92	−2.61	6.32
Ethnic Vote Share (2010)	3.67	−5.97	9.43
Ethnic Vote Share (2014)	1.33	−4.14	6.11
Community			
Centar Sarajevo	37.47	Significance at 0.05 level:	No
Novo Sarajevo	45.34	Significance at 0.05 level:	No
Srebrenik	43.29	Significance at 0.05 level:	No
Tuzla	32.84	Significance at 0.05 level:	No
⋮	⋮	⋮	⋮

Note: Fixed-effects of other communities omitted here. The upper and lower bounds correspond to the 95% confidence interval.

ethnic voting in both pre- and postwar periods.⁷ And except for Srebrenik, the rest of these communities have lower prewar ethnic concentration measured by Herfindahl–Hirschman indices compared to the average of the sample (Hadzic, Carlson, & Tavits, 2020). This implies that in regions without a long-standing tradition of ethnic voting, the mechanism by which wartime violence affects ethnic voting may be different. In areas with a strong culture of cooperation and mild ethnic divisions, it may be easier to return to this norm after the ethnic conflict without a heightened risk of ethnic polarization. This process is less likely or much slower in regions where the culture of cooperation is absent.

These empirical findings from estimating the quantile model help to reconcile the two competing theories of the effect of wartime violence on ethnic voting. This suggests that the concrete effect of violence may be dependent on

⁷ Here, we look at the estimates at the 0.1th quantile. The communities Centar Sarajevo, Novo Sarajevo, Srebrenik, and Tuzla have respectively 52%, 51%, 58%, and 33% prewar ethnic vote shares, and on average 50%, 58%, 54%, and 44% postwar ethnic vote shares, all below the 0.1th quantile (65%) of the proportion of ethnic voting among all communities in the sample. The populations of these communities are, on average, also richer and older than the sample average.

the prior inter-ethnic relations in different communities. Thus, even this simple replication study using quantile models opens up interesting possibilities for further research, which could delve deeper into the local effects that explain the variation of ethnic voting between different regions.

2.4 Example of Political Representation

As another example, political representation by parties is usually not about reflecting average societal preferences. Parties tend to have closer ties to certain societal groups, and more conflictual relations with others. Because these relations can be disrupted or enhanced by external threats, Weschle (2019) argues that economic crises impact the way parties represent the society, which further depends on whether the parties have cooperative or conflictual relations with different societal groups. Under external threats, political parties need not only to be responsive to their own constituencies, but also be responsible to other societal groups in order to unite the whole nation to better cope with the threats (Bueno de Mesquita, 1981; Chowanietz, 2011; Indridason, 2008; Lijphart, 1996). This consequently leads to changes in the parties' societal relations with the groups they are closest to and least close to. More specifically, parties have to trade-off between representing the interests of their own constituencies faithfully and being responsive to other societal groups for a collective effort in handling external threats. Therefore, in times of crises, we would expect empirically that parties' relations with the societal group they are closest to become less cooperative, while their relations with the groups they are least close to become less conflictual (Weschle, 2019). However, these heterogeneous relations are difficult to investigate using the traditional mean-based models, which merely reveal a single average effect.

To investigate the influence of economic crises on the political representation of diverse societal groups, Weschle (2019) developed a latent network model, which draws inferences about the relationship between political parties and societal groups. Subsequently, he utilized quantile models to analyze the effects of economic crises on parties' relationships with different types of societal groups. The latent network model allows for identifying parties' societal relations by estimating the cooperation scores, which measure the level of cooperation or conflict between a party and a societal group for eleven euro-zone countries between 2001 and 2011. Using this measure as the dependent variable, the quantile models estimate varying effects of economic crises on political representation, with different quantiles capturing different parts of the latent cooperation-score distribution. More specifically, higher quantiles in the

model indicate the societal groups with which parties have better initial relations, while lower quantiles focus on those groups that have a less cooperative relationship with the parties.

To demonstrate how quantile models improve our substantive understanding of political representation in times of crises, we replicate the analysis of Weschle (2019). The following codes load the data and draw quantile fitting lines against a scatter plot. Here, the variable `coopscore_1` represents a random draw of the cooperation scores from the latent network model of political representation, and GDP growth per capita is used to examine the impact of economic crises.⁸

```

1  # Load and clean data
2  library(foreign)
3  data <- read.dta("data.dta")
4  dat = data[which(data$type==3),]
5  # Scatter plot and quantile fitting lines
6  ggplot(dat, aes(x = gdppcgrowth, y = coopscore_1)
7    ) +
8    geom_point(color = "gray", alpha = 0.5,
9    size = 2, shape = 1 ) +
10   geom_quantile(quantiles = c(0.01, 0.5, 0.99),
11   aes(linetype = factor(after_stat(quantile))),
12   linewidth = 0.8, color = "black",
13   show.legend = TRUE) +
14   scale_linetype_manual(
15   name = "Quantiles",
16   values = c("dotted", "solid", "dashed"),
17   labels = c("0.01th Quantile", "0.5th Quantile (
18     Median)", "0.99th Quantile")) +
19   labs(x = "GDP Growth per Capita (%)",
20   y = "Cooperation Score") +
21   theme_bw(base_size = 12) +
22   theme(plot.title = element_text(
23     face = "bold", size = 14,
24     hjust = 0.5, margin = margin(b = 10)),
25   axis.title = element_text(face = "bold"),
26   panel.grid.major = element_line(linewidth = 0.2),
27   panel.grid.minor = element_blank(),
28   legend.position = c(0.3, 0.85),
29   legend.background = element_rect(fill = "white",
30     colour = "white"),
31   legend.key.width = unit(1.5, "cm"),
32   plot.caption = element_text(hjust = 0, face = "
33     italic")) +
34   coord_cartesian(ylim = c(-0.15, 0.15)) +

```

⁸ The replication file can be downloaded via the link <https://doi.org/10.7910/DVN/XVPHO1>.

```

31 | annotate(
32 |   "text", x = Inf, y = -Inf,
33 |   label = "Gray circles: Observation points\nLine
      types: Quantile regression estimates",
34 |   hjust = 1.1, vjust = -0.5,
35 |   size = 3, color = "black")

```

Figure 10 shows the three quantile fitting lines in the 0.01th, 0.5th, and 0.99th quantiles that are superimposed onto the scatter plot between GDP growth per capita and cooperation scores. The upper quantile (0.99th quantile) represents societal groups with which a party has a cooperative relationship, while the lower quantile (0.01th quantile) represents those with which a party has a conflictual relationship. The median (0.5th quantile), in contrast, represents groups with which a party has neither a close nor conflictual relation.

According to the fitted quantile lines in Figure 10, there is no discernible effect at the median, but the opposing effects at the lower and upper quantiles of 0.01 and 0.99 are visible. This suggests that the impact of the economic crisis on the relation between a party and a societal group may vary depending on the initial relationship between them. Specifically, as the economy expands, it will enhance parties' cooperative relations with societal groups they are most aligned with (dashed line of the 0.99th quantile), whereas economic crises will




Figure 10 Scatter plot of GDP growth per capita and cooperation scores between political parties and societal groups

damage these relationships, resulting in reduced cooperation. Conversely, as indicated by the dotted fitting line for the 0.01th quantile, during crises, parties endeavor to improve their relations with societal groups they originally have conflictual relations with. These results from the quantile models suggest that economic crises tend to disrupt representation and motivate politicians and parties to “put politics aside” by cooperating more with previously distant groups (Weschle, 2019). If researchers were to rely solely on mean-based models, these new theoretical and empirical perspectives would likely have gone uninvestigated.

Note that the above analysis simply examines the relation between the main explanatory variable and the response variable. To account for potential confounding variables, the next step is to include them in the full model specification. Weschle’s (2019) study includes control variables such as trade openness, population, election year, number of parties, number of events in political communication, and mean cooperation scores for each country and year, which are also included in this replication analysis. To demonstrate the varying effects of economic conditions on political representation, we can specify the lower (0.25th), middle (0.5th), and upper (0.75th) quantiles and estimate the respective conditional quantile effects.⁹ Using the first random draw of cooperation scores, the following codes complete the task and summarize the estimation results for each quantile. Note that compared to the original analysis, the response variable is rescaled by a factor of 1,000 to make the estimates more interpretable.

```

1 # Rescale the response variable by a factor of
  1000
2 dat$coopscore_1_rescale = dat$coopscore_1 * 1000
3 # Specify quantile model
4 mod1 <- rq(coopscore_1_rescale ~ gdppcgrowth +
  openness + population_log + elec + nparties +
  events + coopscore_mean_1, tau = c
  (0.25,0.5,0.75) ,data = dat)
5 # Plot the coefficients
6 plot(mod1)
7 # Summarize the results
8 summary(mod1)

```

Figure 11 demonstrates that the quantile model reveals considerable variation of the effects of different covariates across quantiles compared to the OLS

⁹ Note that in Weschle’s (2019) study, the author specifies the 0.01th, 0.05th, 0.25th, 0.50th, 0.75th, 0.95th, and 0.99th quantiles, and demonstrates in general increasing effects of GDP on political cooperation with larger quantiles. Here, for simplicity, I use only the 0.25th, 0.5th, and 0.75th quantiles.




Figure 11 Coefficient estimates at three conditional quantiles (0.25th, 0.5th, and 0.75th) for political representation example

Note: Black dotted lines represent quantile estimates and horizontal dashed lines represent OLS estimates.

estimates shown by the dashed horizontal lines. For all the control variables, including the intercept, the estimated coefficients have opposite directions between the lower and upper quantiles. The large discrepancy between estimates of the OLS and median regression (quantile model evaluated at 0.5th quantile) also suggests that the heterogeneity in the data cannot be well accounted for by the mean-based model.¹⁰

In addition to the average estimates by the OLS, the results of Table 4 indicate that the effect of GDP growth per capita on political representation varies across quantiles. In the lower quantile (0.25th quantile) of the response variable, the effect is significantly negative, while in the upper quantile (0.75th quantile) the effect is significantly positive. Although the median estimate is statistically distinguishable from zero, its magnitude is substantively very small. This suggests that on average economic conditions have little effect on political representation, but when the economy is doing well, parties are

¹⁰ It also suggests that the restrictive OLS assumptions are not well applied to the data.

Table 4 OLS and quantile estimates for political representation example

Covariates	OLS	0.25th quantile	0.5th quantile	0.75th quantile
(Intercept)	4.057 (2.193)	-4.145* (0.427)	-0.031 (0.035)	2.558* (0.283)
GDP growth pc	0.006 (0.029)	-0.035* (0.002)	-0.002* (0.000)	0.007* (0.002)
Trade openness	0.002 (0.003)	0.003* (0.001)	0.000* (0.000)	0.002* (0.000)
Population (log)	-0.279* (0.131)	0.175* (0.022)	-0.001 (0.002)	-0.127* (0.015)
Election year	-0.276 (0.168)	-0.127* (0.014)	-0.005* (0.002)	0.110* (0.014)
Number of parties	0.081 (0.064)	0.083* (0.008)	0.003* (0.001)	-0.031* (0.005)
Number of events	-0.000 (0.000)	0.000* (0.000)	0.000* (0.000)	0.000* (0.000)
Mean coop. score	2,701.204* (318.297)	-147.077 (79.113)	51.094* (6.866)	413.781* (55.719)

The asterisk * denotes significance at 95% confidence level.

Standard errors are shown in parentheses.

more likely to represent their core constituents. Conversely, when the economy is doing poorly, parties appear to move toward less conflictual relations with groups with which they originally conflict. Thus, this finding indicates that in times of economic crisis, political parties must navigate a distinct balance between catering to their own constituencies and addressing the interests of diverse societal groups to foster national unity. Compared to normal times, the relation between political parties and the societal groups they are least close to becomes less conflictual during periods of crisis. This is the kind of heterogeneous adjustment that a mean-based model is poorly equipped to detect.

As also shown in the table, the coefficients of the rest of the variables, such as population, election year, number of parties, and mean cooperation score, are significantly different between lower and upper quantiles.¹¹ These results

¹¹ Note that the variable “election year” is a dummy variable indicating whether it is the election year in a specific country.

are important for understanding democratic representation in times of crisis, which would remain unknown without exploring the quantile behavior of the explanatory variables over the distribution of the response variable.

To eliminate concerns about potential heterogeneity across countries and over years, we may also control for the country- and/or year-fixed effects. For simplicity, the example presented here uses a single draw of cooperation scores as the response variable. Yet the example is sufficient to demonstrate the substantive payoff of quantile analysis. To have a more complete analysis, more observations can be easily included from the data of Weschle (2019), which will generate similar results.

2.5 Summary

The discussion in this section has shown that, although mean-based models remain central to empirical analysis, they are not always well suited to questions involving substantial heterogeneity. Concentrating on the means has prevented researchers from using more suitable techniques for analyzing heterogeneous political phenomena. For many political studies, the quantile approach allows for the evaluation of heterogeneous data that cannot be satisfactorily analyzed by mean-based models. It not only provides comprehensive summary statistics for the location and spread of a given sample, but also reveals a more detailed relationship between the dependent and explanatory variables. The utility of quantile models is not limited to examining heterogeneous data. Even when the primary interest is not heterogeneity itself, quantile models can still serve as a valuable diagnostic by revealing whether empirical relationships are stable across the distribution.

Using the examples of ethnic voting and political representation, this section illustrates the utility of the quantile approach for applied political research examining continuous response variables in multivariate settings. It shows how quantile models can be beneficial for exploring heterogeneous political units and identifying varying effects across the distribution of the outcomes. The new results have unveiled intriguing insights that were previously undisclosed by the traditional mean-based models.

For instance, in the case of the ethnic voting example, contrary to the average finding indicating a singular positive effect of wartime violence on postwar ethnic voting, the results based on the quantile model demonstrate that the effect is contingent on the degree of prewar ethnic polarization. In communities where there is a strong prewar norm of inter-ethnic cooperation, it is much easier to return to the norm without an increased risk of postwar ethnic polarization. Conversely, in communities with historically high ethnic divisions, the

memory of conflict may instead intensify ethnic polarization. Revisiting the example of political representation also reveals heterogeneous effects of economic crises on political representation of parties with different societal groups. While economic conditions have little impact on political representation on average, parties tend to be more responsive to groups with which they originally conflict when the economy falters. These findings provide crucial insights that would otherwise remain unknown.

Overall, these examples highlight the broader contribution of quantile models to political analysis. They allow researchers to move beyond average effects, to detect substantively meaningful heterogeneity, and to generate more thorough interpretations of political processes. The sections that follow build on this foundation by extending the quantile framework to additional outcome types and modeling settings.

3 Quantile Models for Binary Choices

3.1 Introduction

Besides the continuous response variables introduced in the [previous section](#), many important political outcomes take the form of discrete choices. Such outcomes are central to the study of vote choice, partisan identification, government formation, legislative policymaking, and many other political decisions. For example, researchers might be interested in exploring how conflict among coalition parties affects the adoption of bills with inherently different success rates, or how ideological factors impact coalition formation among coalitions with varying likelihoods of entering government. However, despite the advances in modeling discrete choice data in political science (see, e.g., Alvarez & Glasgow, 1999; Beck, Katz, & Tucker, 1998; Carter & Signorino, 2010; Glasgow, 2011, 2022; McGrath, 2015; Poole, 2000; Rainey, 2016; Sartori, 2003; Traunmüller, Murr, & Gill, 2015), existing estimators are typically centered on conditional means or choice probabilities and pay less attention to distributional heterogeneity. This can lead to incomplete or even inaccurate inferences when heterogeneity among individuals or subgroups of a population is averaged out in the estimation process of discrete response variables.

Because quantiles have useful equivariance properties under monotonic transformations (Koenker & Machado, 1999), they are well suited for modeling discrete choice data which usually require in estimation a certain form of nonlinear transformation (Horowitz, 1992). As with quantile estimation of continuous response variables, quantile models enable a full examination of

the conditional properties for discrete response variables. Compared to the conventional mean-based discrete choice models, quantile models are more robust to certain forms of location-scale shifts of the conditional distribution of the response variable (Benoit & Van den Poel, 2012). Therefore, they are less sensitive to distributional misspecification and provide a more comprehensive view of the effect of the explanatory variables on the response (Benoit, Alhamzawi, & Yu, 2013; Benoit & Van den Poel, 2012; Koenker & Hallock, 2001; Kordas, 2006; Oh, Park, & So, 2016).

This section introduces quantile models for binary choice, the most common form of discrete outcome in political research. Before turning to the binary quantile model, it is useful to consider why conventional mean-based discrete-choice models may be insufficient for the analysis of heterogeneous political phenomena.

3.2 Limitations of Mean-Based Choice Models

To motivate the use of quantile methods for discrete-choice data, it is useful to begin with the logic of conventional binary choice models and to consider their main limitations. For illustration, I focus on the simplest scenario of binary choices where an actor chooses between two alternatives and selects the one that yields the higher utility. The estimation follows the random utility framework, in which observed choices are understood as arising from latent utilities.

Formally, suppose a political actor i faces two choices, $Y \in \{0, 1\}$, and her utilities from choosing the two options are respectively:

$$\begin{aligned} U(y_i = 1) &= \mathbf{x}_i' \boldsymbol{\beta}_1 + \varepsilon_{i1}, \\ U(y_i = 0) &= \mathbf{x}_i' \boldsymbol{\beta}_0 + \varepsilon_{i0}, \end{aligned} \quad (3.1)$$

where x_i is a set of actor-specific characteristics (explanatory variables) of actor i , $\boldsymbol{\beta}_0$ and $\boldsymbol{\beta}_1$ are $k \times 1$ vectors of coefficients with respect to choosing 0 and 1, and $\varepsilon_i = \{\varepsilon_{i1}, \varepsilon_{i0}\}$ includes the error terms. Therefore, the probability of choosing $Y = 1$ is simply the probability of $U_i(Y = 1) > U_i(Y = 0)$, which can be written formally as

$$\begin{aligned} Pr(y_i = 1 | \mathbf{x}_i) &= Pr(U(y_i = 1) > U(y_i = 0)) \\ &= Pr(\mathbf{x}_i'(\boldsymbol{\beta}_1 - \boldsymbol{\beta}_0) + \varepsilon_{i1} - \varepsilon_{i0} > 0) \\ &= Pr(\mathbf{x}_i' \boldsymbol{\beta} + \varepsilon_i > 0), \end{aligned} \quad (3.2)$$

where $\boldsymbol{\beta} = \boldsymbol{\beta}_1 - \boldsymbol{\beta}_0$ and $\varepsilon_i = \varepsilon_{i1} - \varepsilon_{i0}$. Since the constant term in the latent utility specification is unchanged by the choice of switching thresholds, we can

follow the conventional practice and normalize the utility of the second choice to 0. Consequently, we have the following latent utility specification:

$$y_i^* = \mathbf{x}_i' \boldsymbol{\beta} + \varepsilon_i. \quad (3.3)$$

By adding a probability measure over Equation (3.3), it follows that

$$Pr(y_i = 1 | \mathbf{x}_i) = Pr(y_i^* > 0) = Pr(\mathbf{x}_i' \boldsymbol{\beta} + \varepsilon_i > 0) = Pr(\varepsilon_i > -\mathbf{x}_i' \boldsymbol{\beta}). \quad (3.4)$$

Under symmetry of the error distribution, we can obtain the following form

$$Pr(y_i = 1) = Pr(y_i^* > 0) = Pr(\varepsilon_i < \mathbf{x}_i' \boldsymbol{\beta}) = F(\mathbf{x}_i' \boldsymbol{\beta}), \quad (3.5)$$

where $F(\cdot)$ is a cumulative distribution function. In general, the conditional mean-based regression for binary outcomes assumes the following form

$$Pr(y_i = 1 | \mathbf{x}_i) = E(y_i | \mathbf{x}_i) = F(\mathbf{x}_i' \boldsymbol{\beta}). \quad (3.6)$$

This formulation summarizes the relationship through a single conditional index and therefore cannot directly represent heterogeneity across different parts of the latent response distribution (Lu, 2020).

Besides the above limitation, the mean-based discrete choice models can produce systematic prediction errors when the assumption of independence of irrelevant alternatives is violated or when the distribution of the error term is misspecified. These two constraints will be elaborated in the subsequent discussions.

3.2.1 Independence of Irrelevant Alternatives

In broader discrete-choice settings, standard logit models often rely on the independence of irrelevant alternatives (IIA) assumption, which follows from particular assumptions about the stochastic components of utility across alternatives. This assumption implies that when considering any two alternatives, the addition of further alternatives will not alter the probability ratio between them. This can be represented formally by the following equation:

$$\frac{Pr(y_i = 1 | \{0, 1\})}{Pr(y_i = 0 | \{0, 1\})} = \frac{Pr(y_i = 1 | \{0, 1, 2\})}{Pr(y_i = 0 | \{0, 1, 2\})}, \quad (3.7)$$

which means that the probability ratio (or odds ratio) of choosing 1 over 0 given the set of alternatives $\{0, 1\}$ is the same by adding an additional alternative 2 to the original set of alternatives $\{0, 1\}$.

The IIA assumption becomes stringent in cases where, for instance, alternatives 0 and 2 exhibit similar characteristics while alternative 1 stands out. In such situations, the probability of selecting 0 decreases due to the presence

of a comparable alternative 2, but the probability of choosing 1 remains relatively unchanged. Consequently, the probability ratio between 1 and 0 increases because of the existence of alternative 2, thus violating the IIA assumption.

As a more concrete example, consider vote choices in multi-party systems. When examining voting behavior in such systems, the emergence of a third party with an ideological position closely aligned with one of the two existing parties will inevitably impact voters' choices among the three parties. Specifically, the presence of the new party is likely to decrease the election chances of the party that shares a similar ideology.

When the IIA assumption is violated, conventional mean-based discrete choice models like logit and probit may lead to biased inferences. While the violation of the IIA assumption is less consequential in binary choice settings, it becomes a problem with the presence of multiple alternatives, which will be examined in Section 4.

3.2.2 *Potential Distributional Misspecification of Mean-Based Binary Models*

The most commonly used binary models, such as logit and probit models, are restricted by their relatively rigid distributional assumptions. To convert a linear combination of covariates with complete support over the real line to a probability space ranging from zero to one, traditional binary models use a certain type of cumulative distribution function as the link function. The most common choices of the cumulative distribution function are the normal cumulative distribution, which gives rise to the probit model:

$$Pr(y_i = 1 | \mathbf{x}_i) = \int_{-\infty}^{\mathbf{x}_i' \boldsymbol{\beta}} \phi(t) dt = \Phi(\mathbf{x}_i' \boldsymbol{\beta}), \quad (3.8)$$

and the logistic cumulative distribution, which is the core of the logit model:

$$Pr(y_i = 1 | \mathbf{x}_i) = \frac{\exp(\mathbf{x}_i' \boldsymbol{\beta})}{1 + \exp(\mathbf{x}_i' \boldsymbol{\beta})}. \quad (3.9)$$

However, these distributional assumptions are substantively stringent and may lead to inaccurate estimations. To demonstrate potential bias that can be generated when the underlying error distribution differs from the assumed one, we can simulate a simple dataset in the following form:

$$y_i^* = \beta_0 + \mathbf{x}_i' \boldsymbol{\beta}_1 + \varepsilon_i$$

$$Pr(y_i = 1 | \mathbf{x}_i) = 1 - F(-\beta_0 - \mathbf{x}_i' \boldsymbol{\beta}_1) \quad (3.10)$$

where $\beta_0 = 2$, $\beta_1 = -1$, $\mathbf{x} \sim \text{uniform}(0, 5)$, and $F(\cdot)$ are the cumulative distributions with the error term assuming respectively the following seven distributions (Lu, 2020):

- **Distribution 1:** Normal distribution: $N(0, 1)$
- **Distribution 2:** T distribution with 3 degrees of freedom
- **Distribution 3:** χ^2 distribution with 3 degrees of freedom
- **Distribution 4:** Asymmetric Laplace distribution (location 0 and scale 1)
- **Distribution 5:** Kurtotic distribution from a weighted mixture of normal distributions $\frac{2}{3}N(0, 1) + \frac{1}{3}N(0, 1/100)$
- **Distribution 6:** Bimodal distribution from a symmetric mixture of normal distributions $(N(-1, 4/9) + N(1, 4/9))/2$
- **Distribution 7:** Skewed distribution with a mixture of three normal distributions $(N(-22/25, 1) + N(-49/125, 9/4) + 3N(29/250, 25/81))/5$

Estimating the data with the above distributions by the probit model, Figure 12 shows that except for the correctly specified model in the top-left panel (distribution 1), the probit model can deviate noticeably from the true response probabilities under alternative latent error distributions. For example, for the estimation of the data with the skewed distribution of the error term, the




Figure 12 Comparison between the predicted probabilities and the true probabilities under different error distributions

probit model under-predicts the probabilities in lower values of X . However, it is more likely to overpredict the probabilities when the error term follows the bimodal distribution. This indicates that in the discrete choice settings, the specification of error terms matters. Once the underlying distribution deviates from the assumption, the estimator is likely to produce biased estimates.

Compared to the mean-based discrete choice models, quantile models adopt more flexible distributional assumptions and are thus more robust to distributional misspecifications. For the binary choices, the following section will introduce the technical details of the binary quantile model.

3.3 Binary Quantile Model

Mean-based discrete choice models, including the logit and probit models, are commonly used to analyze data with discrete outcomes. However, as demonstrated earlier, these conventional mean-based models impose strong assumptions about data distributions and are likely to overlook heterogeneity among individuals or subgroups of a population. In many research situations, we are interested in how explanatory variables operate across the latent response distribution beyond just the conditional mean.

Quantile models represent a particularly useful approach for this purpose. For a binary response variable, we assume that a latent continuous variable Y^* determines the observed binary outcome Y . Therefore, a coefficient $\beta_{j,\tau}$ from the quantile estimation represents the change in the τ th quantile of the latent propensity Y^* for a one-unit increase in predictor X_j , holding other variables constant. Coefficients at higher quantiles reflect effects on observations with high latent propensity, while coefficients at lower quantiles reflect those with low latent propensity. Thus, unlike mean-based binary models, the binary quantile model captures heterogeneous effects across the latent propensity distribution.

For binary outcomes, the quantile-regression logic cannot be applied directly to the observed response in the same way as in the continuous case, and estimation therefore requires a latent-variable formulation together with simulation-based methods. To handle the problem, scholars have begun to explore the advantages of the Bayesian approach for quantile estimation (Benoit, Alhamzawi, & Yu, 2013; Benoit & Van den Poel, 2012; Florios & Skouras, 2008; Kordas, 2006). In particular, Benoit and Van den Poel (2012) developed a simulation-based estimator for binary quantile models that draws posterior samples through the MCMC process. This approach allows inference of parameter values from the joint posterior distribution, without the need to adopt ad hoc approximation methods.

Because the minimizing problem in Equation (2.5) is closely related to the asymmetric Laplace distribution (ALD) (Koenker & Machado, 1999), it is possible to estimate the quantile regression within the framework of the ALD distribution. The ALD density of a random variable y conditioning on three parameters, μ , σ , and τ , can be written as

$$f(y|\mu, \sigma, \tau) = \frac{\tau(1-\tau)}{\sigma} \exp \left\{ -\rho_{\tau} \left(\frac{y-\mu}{\sigma} \right) \right\}, \quad (3.11)$$

where $\rho_{\tau}(x) = x(\tau - I(x < 0))$ is determined by the specified quantile, and $\mu = \mathbf{x}'\boldsymbol{\beta}_{\tau}$ containing the covariates is the main parameter of interest (Yu & Zhang, 2005). Compared to other commonly used distributions for mean-based discrete choice models, the ALD is more flexible in the sense that it can easily adjust the skewness, shape, and location by varying the values of the parameters. In a discrete choice setting, the dispersion parameter σ is normalized to 1 for identification.

By inserting a monotone indicator function, the quantile representation of the binary response variable given a set of covariates has the following form:

$$Q_{\tau}(y_i = 1|\mathbf{x}_i) = I(F_{\tau}^{-1}(\mathbf{x}_i'\boldsymbol{\beta}_{\tau}) > 0), \quad (3.12)$$

where $I(\cdot)$ is an indicator function that takes the value of one when the condition within the bracket is satisfied, and zero otherwise. For a binary response variable y_i , the probability of $y_i = 1$ conditional on the data and the quantile is as follows:

$$\begin{aligned} Pr(y_i = 1|\mathbf{x}_i, \boldsymbol{\beta}_{\tau}, \tau) &= Pr(y_i^* > 0|\mathbf{x}_i, \boldsymbol{\beta}_{\tau}, \tau) \\ &= Pr(\mathbf{x}_i'\boldsymbol{\beta}_{\tau} + \varepsilon_i > 0|\mathbf{x}_i, \boldsymbol{\beta}_{\tau}, \tau) \\ &= 1 - F_{\text{ALD}}(-\mathbf{x}_i'\boldsymbol{\beta}_{\tau}|\mathbf{x}_i, \boldsymbol{\beta}_{\tau}, \tau). \end{aligned} \quad (3.13)$$

Instead of using Metropolis–Hasting within the Gibbs algorithm to sample the joint posterior distribution, the ALD can be represented as a location-scale mixture of normal distributions (Kotz, Kozubowski, & Podgorski, 2001, 2003; Kozumi & Kobayashi, 2011; Reed & Yu, 2009; Yu & Moyeed, 2001):

$$\varepsilon = \theta z + \tau\sqrt{z}u, \quad (3.14)$$

where

$$\theta = \frac{1-2p}{p(1-p)}, \text{ and } \tau = \sqrt{\frac{2}{p(1-p)}}. \quad (3.15)$$

Here, p denotes the skewness of the distribution. As a result, latent variable y_i^* can be rewritten as

$$y_i^* = \mathbf{x}_i'\boldsymbol{\beta}_{\tau} + \theta z_i + \tau\sqrt{z_i}u_i, \quad (3.16)$$

where $z_i \sim \text{Exponential}(1)$ and $u_i \sim \text{Normal}(0, 1)$ are auxiliary variables. Consequently, we have the joint distribution given data as

$$f(\mathbf{y}^* | \boldsymbol{\beta}_\tau, \mathbf{z}) \propto \left(\prod_{i=1}^N z_i^{-1/2} \right) \exp \left\{ - \sum_{i=1}^N \frac{(y_i^* - \mathbf{x}_i' \boldsymbol{\beta}_\tau - \theta z_i)^2}{2\tau^2 z_i} \right\}. \quad (3.17)$$

In the simple binary choice model, it is possible to augment the data in the sampling procedure by adding a latent variable y_i^* , which has continuous support over the whole real line:

$$\pi(y_i^* | y_i, \boldsymbol{\beta}_\tau, \tau) \propto \{I(y_i^* > 0)I(y_i = 1) + I(y_i^* < 0)I(y_i = 0)\} f_{\text{ALD}}(y_i^* | \mathbf{x}_i' \boldsymbol{\beta}_\tau, \tau). \quad (3.18)$$

As a result, the inference of $\boldsymbol{\beta}_\tau$ is simply the integral of latent variable y_i^* :

$$Pr(\boldsymbol{\beta}_\tau | y_i) = \int_{y_i^*} p(\boldsymbol{\beta}_\tau | y_i, y_i^*) f(y_i^* | y_i) dy_i^*. \quad (3.19)$$

The estimation procedure following the above calculations has been incorporated into an R package *bayesQR* (Benoit & Van den Poel, 2017). Thus, rather than dwelling on the details of the sampling methods, the following section will demonstrate the application of the binary quantile model using a real-world political example on the EU policymaking.

3.4 Example of Binary Choices: EU Policymaking

To illustrate the application of the binary quantile model, this section considers an example of policymaking in the European Union. In 1992, the Maastricht Treaty established the so-called codecision legislative procedure, which requires the Council of the European Union to negotiate with the European Parliament (EP) in order to pass a bill. While the introduction of the codecision procedure empowered the EP to enhance the democratic process, the Amsterdam treaty of 1999 amended the procedure to allow legislation to be concluded as early as in the first reading stage. This reform raised trade-offs between democracy and efficiency, and gave the colegislators the option of concluding legislation earlier or later in the process. The potential for concluding a bill early in the EU legislative procedure has led to more frequent informal meetings and sparked the concern that a biased rapporteur, who acts as the EP's representative, might exploit this opportunity to advance personal policy goals at the expense of the whole EP (Farrell & Héritier, 2004).

To study the conditions under which colegislators, that is, the Council and the EP, decide to conclude an early agreement, Rasmussen (2011) proposes three sets of factors that may influence early conclusion of EU legislative bills, namely, the delegation costs of the rapporteurs from the EP, the bargaining

certainty, and the bargaining impatience. While the first factor is expected to reduce the likelihood of a first reading agreement, the latter two factors are hypothesized to increase the likelihood. The empirical results from the ordinary logit regression lend support to the three proposed hypotheses.

However, the ordinary logit model originally used by the author is unable to account for the potential heterogeneity in the EU codecision bills. For example, the impact of delegation costs on early agreement may differ depending on each bill's latent propensity to be concluded through a fast-track procedure. For conflictual bills that are less likely to be included in fast-track legislation or trivial bills where consensus among legislators is easy to reach, it is reasonable to expect that the impact of delegation costs on early agreement will be low (Lu, 2020). Yet, for bills that have neither a very low nor a very high likelihood of being concluded early in time, we would expect a more significant impact of delegation costs.

To demonstrate the new insight that could possibly be generated from the quantile approach, I replicate the original analysis in this section by applying the binary quantile model (see also Lu, 2020). In the model, the conditional quantiles represent varying latent propensity of early agreement for different bills. Although the IIA assumption is not relevant in this case as there are only two choice alternatives (early conclusion of a bill or not), the key issue lies in the identification of unobserved heterogeneity among legislative bills. Even by controlling for potential confounding variables in the mean-based model, it still provides no guarantee that unobserved heterogeneity among bills and their local effects has been well accounted for. To identify unobserved heterogeneity and promote our understanding of the lawmaking in the EU, I focus on the full model (Model III in table 2 and figure 4 of the original paper by Rasmussen (2011)) and estimate the data using the binary quantile model, with specified quantiles ranging from 0.1 to 0.9.

Using a cleaned version of the original dataset, the following codes load the ready-for-estimation data into the memory and carry out a brief preliminary inspection.¹²

¹² The data are publicly available and can be downloaded through the link https://journals.sagepub.com/doi/suppl/10.1177/1465116510388675/suppl_file/10.11771465116510388675-replication_files.zip. The correspondence between variable names and their meanings is as follows: choice: Early agreement (dummy); var6: Policy distance rapporteur-EP median; var3: Policy distance rapporteur-EP median * Big party group; var5: Big party group; var10: Country coherence of key negotiators; var15: Closeness of working relationship (calendar year); urgcy_yes: Urgency of legislation: existing legislation expires; urgcy_noleg: No existing legislation; var4: Workload (pending legislation during first reading); var11: Issue areas consulted (no. of consultative committees); var14: New act; var7: Directive; var17: Inter-institutional disagreement (Rasmussen, 2011).

```

1 # Load EU Legislative data
2 load("data_legislature.RData")
3 # Check variable names
4 names(data_legislature)
5 # Have a look at the first 5 rows of the
   datasets.
6 head(data_legislature,5)

```

To implement the simple binary quantile estimation, we can use the `bayesQR` function provided by the package of the same name.¹³ The following codes demonstrate the estimation procedure, where the estimation results are stored in the object `mod1`, which can be further assessed using the function `summary`. It is also possible to predict the responses relying on the estimated coefficients using the function `predict`.

```

1 # Load the package
2 library(bayesQR)
3 # Estimate the model
4 mod1 <- bayesQR(choice ~ var6 + var3 + var5 +
   var10 + var15 + urgcy_yes + urgcy_noleg + var4
   + var11 + var14 + var7 + var17,
5 data = data_legislature, quantile = c
   (0.1,0.25,0.5,0.75,0.9),ndraw = 10000)
6 # Summarize the results
7 mod1_out <- summary(mod1)

```

As the Bayesian procedure is used for the estimation of the binary quantile model, we have to check whether the MCMC chains of the estimates have converged before proceeding to the interpretation of the results. This can be done by creating traceplots with the following functions.

```

1 # Traceplot of MCMC draws
2 par(mfrow = c(2,3), mar = c(4, 4, 1, 2))
3 plot(mod1, var = "var3", plottype = "trace",
   burnin = 100,main = "",ylab = "Coefficient",
   xlab = "Iteration")

```

Figure 13 suggests satisfactory mixing and no obvious convergence problems. Consequently, we proceed with the substantive interpretation of the estimates.

To examine, for instance, the estimated coefficients of the distance between the rapporteur and the EP median (i.e., the delegation costs), Figure 14 presents the quantile coefficient plot. While the logit model indicates a significantly

¹³ Note that alternatively we can also use the `cbq` function from the `cbq` package for the estimation of binary quantile regression.




Figure 13 Traceplot of MCMC draws




Figure 14 Coefficients of distance between rapporteur and EP median

Note: The black dotted line and the shaded area represent the quantile estimates and their 95% credible intervals, while the solid lines represent the estimates by the logit model with 95% confidence interval.

negative average estimate, the quantile models reveal different effects at the lower and upper quantiles of the latent propensity for early agreement. More specifically, the figure shows that, although the median (0.5th quantile) effect mirrors the significantly negative mean estimate, there are no significant effects at the lower (0.1th quantile) and upper (0.9th quantile) ends of the latent early agreement propensity.

These estimates suggest that compared to an average bill, the effects of the subgroups of bills at the tails of the distribution of the response variable can be quite different. Substantively, it indicates that for contested bills that are unlikely to be concluded early in the EU legislative process, or trivial technical bills that are mostly decided by technocrats behind closed doors, the impact of potential policy deviation of negotiators on early conclusion can be different compared to other ordinary bills. In other words, the negative impact of biased policy negotiators on early conclusion of a bill only holds if the bill under consideration is neither very likely nor very unlikely to be considered as a fast-track legislation for early agreement. Therefore, the concern about the policy influence of a biased rapporteur in the fast-track legislation is only partially eliminated for those average bills, but not for bills at the tails of the distribution. The quantile finding also implies that the inclusion of bills in the fast-track procedure is a strategic decision of the EU legislators depending on the nature of the bills, and thus provides new insight – that would have remained unexplored using the conditional mean-based model – for understanding the legislative activities in the EU.

For the comparison of model performances between the ordinary logit model and binary quantile models, we can predict probabilities based on the estimates, and compare the predicted choices with the observed outcomes.

```
1 predict(mod1,X = data_legislature,burnin = 100)
```

The results show that compared to the 76.6% prediction accuracy of the ordinary logit model, the prediction accuracies of the binary quantile model in nine quantiles range from 75.1% to 77.7%. Five out of nine quantile predictions perform equal to or better than the ordinary logit model. More importantly, however, the substantive gain of the quantile approach lies not primarily in prediction, but in its ability to uncover heterogeneity in the relationship between covariates and early agreement.

To further examine the effects of other variables, we can also plot for all covariates the coefficient estimates and their uncertainty (Lu, 2020). In essence, if unobserved heterogeneity has been well controlled for, we should expect no large variation among different quantile estimates. However, as shown in Figure 15, the variation is large for the quantile estimates of many variables that have been included in the analysis. This indicates that compared to the ordinary logit model, quantile models are more useful for the examination of the heterogeneous relationship between those covariates and the response variable.

If we look more closely into the details of the estimation results, we can find that the predicted probabilities at quantiles 0.1 and 0.9 have the smallest reductions when delegation costs are measured by the distance between the




Figure 15 Quantile coefficient plot for all variables in EU policymaking example

rapporteur and the EP median. This reinforces the interpretation that the EU colegislators are less concerned about delegation costs when the bills are very likely or unlikely to be concluded early in time. Such bills, for example, include those that are of an administrative nature and thus need no further negotiation, or those on which it is difficult for member states to reach a compromise, and consequently, they reach no agreement at all. These are significant insights that emerge from estimating quantile models, which are likely to be overlooked if one relies solely on mean-based models.

3.5 Summary

Discrete response variables, such as the presence of conflicts, the voting behavior of electorates, and the success or failure of legislative bills, are often central

to empirical political research. Despite the common occurrence of discrete choice data, most applications of quantile methods concentrate on continuous response variables, leaving the discrete case comparatively less explored. This section has shown that the quantile perspective can also be extended to binary choice settings, where it provides a useful complement to conventional mean-based models. Rather than summarizing behavior solely through conditional choice probabilities, the binary quantile model allows researchers to examine heterogeneity across the latent response distribution underlying observed choices.

Focusing only on average effects can provide an incomplete account of political behavior when the underlying process is heterogeneous. In binary settings, this problem does not disappear simply because the observed outcome takes values of zero and one. What matters is that the latent propensity toward the outcome may vary systematically across observations, and that explanatory variables may affect different parts of that latent distribution in different ways. Binary quantile models are valuable because they make such variation visible.

From the EU legislative example, it is evident that compared to an average bill, the EU legislators are less concerned about the delegation costs of bills that are either very likely or very unlikely to be passed early. These bills typically pertain to technical matters or involve contentious political issues. Thus, even in the simple two-alternative cases, unobserved heterogeneity justifies the use of quantile models. These models effectively capture heterogeneity in the effects of main explanatory variables, therefore providing new insights on the EU lawmaking process.

The application of quantile models for discrete choice data is not limited to binary cases. In the [following section](#), it extends the discussion to quantile models for analyzing multiple alternatives, and for cases where there might also be alternative specific-features in choice sets.

4 Quantile Models for Multiple Choices

4.1 Introduction

The simple binary choice framework discussed in the [previous section](#) is useful for simple two-alternative settings, but many political decisions involve more complex choice environments. In political research, for example, we frequently analyze observational datasets where decision-makers operate under heterogeneous decision-making contexts, characterized not only by distinct option menus across cases but also by intra-case variability in available alternatives.

In such settings, a model designed for simple binary outcomes is no longer sufficient. A useful way to address this problem is to reformulate multi-alternative choice as a set of conditional binary comparisons within each choice set. This is the logic behind the conditional binary quantile (CBQ) model, which extends quantile analysis to settings with multiple alternatives and alternative-specific features (Lu, 2020).

In the context of binary choices, the conditional probability that a specific alternative is selected from a choice set can be represented as the likelihood that the latent utility of this alternative surpasses that of the other option. However, with the increasing number of choice alternatives, the probability calculation becomes more complicated. Consequently, additional statistical procedures are necessary to incorporate multiple alternatives and account for alternative-specific features, which will be introduced in the following sections.

4.2 Incorporation of Multiple Alternatives and Alternative-Specific Features

Consider the following example with three alternatives in a choice set $J = \{1, 2, 3\}$. The probability of choosing $j = 1$ given a set of covariates $\mathbf{x}_i$ is the probability that the utility from alternative 1 is greater than those of alternatives 2 and 3:

$$Pr(y_{i1} = 1|\mathbf{x}_i) = Pr(y_{i1}^* > y_{i2}^* \quad \text{and} \quad y_{i1}^* > y_{i3}^*|\mathbf{x}_i), \quad (4.1)$$

where $\mathbf{x}_i = (\mathbf{x}_{i1}, \mathbf{x}_{i2}, \mathbf{x}_{i3})'$ collects the alternative-specific covariates for all three options in choice set i (Lu, 2020). Modeling the random component in a latent linear specification $y_{ij}^* = \mathbf{x}_{ij}'\boldsymbol{\beta} + \varepsilon_{ij}$, we can rearrange the equation and rewrite the conditional probability as follows:

$$\begin{aligned} Pr(y_{i1} = 1|\mathbf{x}_i) &= \int_{-\infty}^{(\mathbf{x}_{i1}-\mathbf{x}_{i2})'\boldsymbol{\beta}} \int_{-\infty}^{(\mathbf{x}_{i1}-\mathbf{x}_{i3})'\boldsymbol{\beta}} f(\varepsilon_{i3} - \varepsilon_{i1}, \varepsilon_{i2} - \varepsilon_{i1}) \\ &\quad \times d(\varepsilon_{i3} - \varepsilon_{i1})d(\varepsilon_{i2} - \varepsilon_{i1}), \end{aligned} \quad (4.2)$$

where $f(\cdot)$ denotes the joint probability density function of $\varepsilon_{i3} - \varepsilon_{i1}$ and $\varepsilon_{i2} - \varepsilon_{i1}$ (Lu, 2020).

We can simplify the above equation by modeling the difference between two random components as a single random variable: $\eta_i^{jk} = \varepsilon_{ij} - \varepsilon_{ik}$. More generally, let u_i^{jk} denote $(\mathbf{x}_{ij} - \mathbf{x}_{ik})'\boldsymbol{\beta}$; the probability of choosing alternative $j \in J$ in the choice set with the number of alternatives K_i can be expressed as an integral over the joint distribution of all pairwise error differences relative to j (Lu, 2020):

$$Pr(y_{ij} = 1 | \mathbf{u}_i) = \int_{-\infty}^{u_i^{j1}} \cdots \int_{-\infty}^{u_i^{j(j-1)}} \int_{-\infty}^{u_i^{j(j+1)}} \cdots \int_{-\infty}^{u_i^{jK_i}} f(\eta_i^{j1}, \dots, \eta_i^{j(j-1)}, \eta_i^{j(j+1)}, \dots, \eta_i^{jK_i}) d\eta_i^{j1} \dots d\eta_i^{jK_i}. \quad (4.3)$$

This representation makes clear that, once more than two alternatives are present, the choice probability depends on a joint comparison across all available options rather than on a single binary threshold. To formulate an appropriate likelihood function, a multinomial structure can be embedded in the calculation of the joint probability (McFadden, 1974). To do so, the probability for each alternative is weighted by the number of choice alternatives in the corresponding choice set. Each weight is formulated as $\frac{K_i!}{\prod_{l \in \{1 \dots S_i\}} s_{il}!}$, where K_i is the number of alternatives in choice set i , and s_{il} is the number of alternatives l in choice set i with the total number of alternatives S_i . This leads to the following conditional likelihood function given N choice sets (Lu, 2020):

$$\mathcal{L}_{\text{condition}} = \prod_{i=1}^N \frac{K_i!}{\prod_{l \in \{1 \dots S_i\}} s_{il}!} \prod_{j=1}^{K_i} Pr(y_{ij} = 1 | \mathbf{x}_i)^{I(y_{ij}=1)} Pr(y_{ij} = 0 | \mathbf{x}_i)^{I(y_{ij}=0)}. \quad (4.4)$$

Equation (4.4) is a general form of the conditional binary choice model. This general framework encompasses the CL model (McFadden, 1974), the conditional probit model (Hausman & Wise, 1978), and the mixed logit (MXL) model (Hensher & Greene, 2003). However, in contrast to these models, the CBQ model relies on less restrictive distributional assumptions and does not depend on the IIA assumption. Compared to the conditional probit model, the CBQ model offers a comparatively parsimonious way to study heterogeneity across the latent utility distribution.

To estimate the CBQ model, the data augmentation procedure as introduced in the simple binary setting cannot be directly applied due to multiple alternatives in each choice set. As a result, there is no clear-cut threshold for the latent utility function. To deal with the problem, the estimation proceeds via the joint posterior density of all random components in the model, which can be formulated as

$$\begin{aligned} & Pr(\boldsymbol{\beta}, \mathbf{y}^* | \mathbf{y}, \tau) \\ & \propto Pr(\boldsymbol{\beta}) \prod_{i=1}^N \frac{K_i!}{\prod_{l \in \{1 \dots S_i\}} s_{il}!} \prod_{j=1}^{K_i} \left(\prod_{k \in \{1 \dots K_i\} / j} F(y_{ij}^* | (\mathbf{x}_{ij} - \mathbf{x}_{ik})' \boldsymbol{\beta}, \tau) \right)^{I(y_{ij}=1)} \\ & \times \left(1 - \prod_{k \in \{1 \dots K_i\} / j} F(y_{ij}^* | (\mathbf{x}_{ij} - \mathbf{x}_{ik})' \boldsymbol{\beta}, \tau) \right)^{I(y_{ij}=0)}, \end{aligned} \quad (4.5)$$

where $F(\cdot)$ is the cumulative distribution function of the ALD, N is the number of choice sets, K_i is the number of alternatives in the choice set i , and s_{il} is the number of alternatives l in the choice set i , with the total number of alternatives S_i (Lu, 2020). The posterior distribution can then be sampled using a gradient-based MCMC algorithm (Hamiltonian Monte Carlo) that provides an efficient sampling routine (Hoffman & Gelman, 2014).

For applied research, there is already a statistical software that wraps up the above estimation equations into an easy-to-use R package (Lu, 2020). The corresponding package for estimating the CBQ model, `cbq` is available in the CRAN, which can be downloaded by executing in the R console the following line of code:

```
1 install.packages("cbq")
2 # Load the package
3 library(cbq)
```

Note that when choice sets contain more than two choice alternatives, we need to specify the conditionality of the estimation by using choice-set indicators. Each choice set must have a unique indicator so that the estimator can identify the alternatives belonging to the same choice set and how they are correlated with each other.

In the package, the conditionality of the estimation on the choice-set indicators is formulated by a vertical bar immediately following the specification of the covariates: “ $y \sim x \mid \text{indicator variable}$.” The model specification in the function “`cbq`” has the following patterns: While we can specify formula in the traditional form of “ $y \sim x$ ” for simple binary choices, for multiple choices with or without varying choice alternatives, choice-set indicators need to be supplied in addition to the above formula: “ $y \sim x \mid \text{choice-set indicator}$.” Moreover, we can estimate fixed- or random-effects in the following form: “ $y \sim x \mid \text{choice-set indicator} \mid \text{fixed-effects} \mid \text{random-effects}$.”

In the following, I will present two prominent real-world political examples to demonstrate the basic elements of the quantile estimation procedure for multiple choices. The applications will also show how to view and interpret the results in multiple-choice and varying-choice-alternatives settings.

4.3 Example of Multiple Choices: US Presidential Election in 1992

The first example concerns the US presidential election in 1992, a natural example of multiple choice. Compared to the simple binary choice setting, there are three alternatives in this example, namely Clinton, Bush, and Perot. Using a subsample of 909 respondents from the 1992 American National Election Study (Miller, Kinder, & Rosenstone, 1993), Alvarez and Nagler (1995) offer

a thorough examination of the mainstream explanations for Clinton's success, including the economic voting, the spatial voting, and the so-called "angry voting" hypotheses.¹⁴ The results from their multinomial probit model show that economic perception is the dominant factor determining the election outcome. While issue and ideological distances between the electorate and the candidates have some explanatory power, they are unlikely to alter the result. In addition, the authors find that the "angry voting" hypothesis was not fully supported by the data.

In contrast to the ordinary logit model, the multinomial probit model employed in the analysis by Alvarez and Nagler (1995) permits simultaneous examination of multiple choices. It also provides a correlation structure between random components of the utility function, thereby eliminating the restrictive IIA assumption (Hausman & Wise, 1978). This is particularly relevant in this example, as the presence of the third choice alternative may affect the voters' utility of choosing other candidates, which would not be taken into account if the IIA assumption were to hold. However, the correlation structure included in the multinomial probit model only distinguishes preferences between choice alternatives and does not differentiate the heterogeneous preferences of individuals for each choice alternative. Thus, it leaves substantial room for the application of the quantile approach in this example.

To revisit the election from a quantile perspective, we can specify varying quantiles from 0.1 to 0.9 as in Lu (2020), which correspond to different points in the latent propensity distribution for choosing a given candidate. Here, the conditional quantiles represent varying latent propensity of voting for a specific candidate given a set of explanatory variables. Consequently, we can estimate effects of the covariates for each specified quantile. As an example, in the following codes, we estimate the 0.1th conditional quantile using the `cbq` function, with other options regarding the details of the MCMC sampling procedure set to default.¹⁵

¹⁴ In this subsample used for estimation, Clinton received 45.8% share of votes, Bush 34.1% and Perot 20.0%, which is close to real outcomes.

¹⁵ The correspondence between the variable names and their meanings is as follows: 1. choice: (1) Bush, (2) Clinton, (3) Perot; 2. educ: Respondents education; 3. east: Region (East); 4. south: Region (South); 5. west: Region (West); 6. women: Gender (Female); 7. respfnp: Felt personal finance were worse; 8. natlec: Felt national economy was worse; 9. bclibdis: Ideological Distance; 10. glibdis: Ideological Distance; 11. rplibdis: Ideological Distance; 12. dem: Democrat; 13. rep: Republican; 14. respjob: Oppose government jobs; 15. resphlth: Oppose government health care; 16. respblk: Oppose government minority assistance; 17. respab: Abortion; 18. termlim: Term limits; 19. age1829: Age: 18-29; 20. age3044: Age: 30-44; 21. age4559: Age: 45-59; 22. newvoter: New or returning voter; 23. deficit: Felt deficit was a major problem; 24. chid: Choice set indicator.

```

1 # Estimate the model
2 mod_vote1 <- cbq(choice ~ dist + respfinp.bush +
3   natlec.bush + respgjob.bush + resphlth.bush +
4   respblk.bush + respab.bush + east.bush + south.
5     bush + west.bush + newvoter.bush + termlim.
6     bush +
7     deficit.bush + dem.bush + rep.bush + women.bush +
8     educ.bush + age1829.bush + age3044.bush +
9     age4559.bush
10    + respfinp.clinton + natlec.clinton + respgjob.
11      clinton + resphlth.clinton +
12      respblk.clinton + respab.clinton + east.clinton +
13      south.clinton + west.clinton + newvoter.
14      clinton + termlim.clinton +
15      deficit.clinton + dem.clinton + rep.clinton +
16      women.clinton + educ.clinton + age1829.clinton
17      + age3044.clinton + age4559.clinton|chid,
18 data = data_vote, q = 0.1)

```

In the computational codes, the parameter “`.|chid`” specifies the choice-set identifier. We can inspect the fitted object stored in `mod_vote1` using the following lines of codes.

```

1 # Print the results
2 print(mod_vote1)
3 # Examine the coefficient estimates
4 coef(mod_vote1)
5 # Check convergence
6 plot(mod_vote1)
7 # Produce coefficient plot
8 plot(mod_vote1, "coef")

```

By making a slight modification to the codes, it is straightforward to estimate more conditional quantiles of the data. In this analysis, I follow the usual practice by specifying nine equally spaced quantiles from 0.1 to 0.9.¹⁶

To examine the performance of the models, the `predict` function can be used to calculate the predicted outcomes, and then compare them to the real observed values. Table 5 illustrates the confusion matrices generated respectively by the multinomial probit and the CBQ model at the 0.1th quantile. The values on the diagonal show the number of times the respective models correctly predicted the true election vote choices, and those off the diagonal show the number of incorrect predictions. While the predictions for the vote choices

¹⁶ The codes for the specification of other quantiles are available in the replication files that come with this Element.

Table 5 Confusion matrices of multinomial probit and CBQ models

		Multinomial probit			CBQ at 0.1th quantile		
	Predicted	Bush	Clinton	Perot	Bush	Clinton	Perot
Actual	Bush	278	31	1	257	22	31
	Clinton	29	385	2	19	368	29
	Perot	87	93	3	63	69	51

of Bush and Clinton are relatively similar between the two models, the prediction by the multinomial probit model for the choice of Perot is much worse than the prediction of the CBQ model at the 0.1th quantile. This suggests that the multinomial probit model appears to underpredict the voting probabilities for Perot at the lower quantiles, which are better estimated by the quantile model.

We can further compare the performance between the CBQ model and the multinomial probit model in the nine specified quantiles using the following codes.

```
1 # Predicted probability of choices at 0.1th
  quantile
2 predict(mod_vote1)
3 # Real choices
4 ids <- data_vote$chid
5 real_choice <- rep(0,909)
6 for (i in 1:909){
7   nnn <- data_vote$choice[which(ids==i)]
8   real_choice[i] <- which.max(nnn)
9 }
10 # Calculate prediction accuracy
11 tmp <- predict(mod_vote1)[,1]
12 pred1 <- rep(0,909)
13 for (i in 1:909){
14   nnn <- tmp[which(ids==i)]
15   pred1[i] <- which.max(nnn)
16 }
17 # Confusion table
18 table(real_choice, pred1)
19 # Prediction accuracy
20 length(which(real_choice == pred1))/909
```

Table 6 reveals that the prediction accuracies of the CBQ model at nine quantiles are all better than the prediction accuracy of the multinomial probit.

Besides a better prediction performance, the CBQ estimates show considerable variation across quantiles. To quantify the unobserved heterogeneity

Table 6 Prediction accuracy of multinomial probit and CBQ models (%)

Multi. logit	CBQ								
	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9
73.3	74.4	75.1	75.4	75.5	75.1	75.5	75.6	75.4	75.5

Table 7 Voter heterogeneity

Covariates	Alternatives	
	Bush	Clinton
Abortion	0.280	0.032
Democrat	2.657	0.512
Republican	0.779	2.723
Respondents' education	0.096	0.019

Heterogeneity is measured by the standard deviation of quantile estimates for each covariate.

among voters for each choice alternative, we can calculate the standard deviation of nine quantile estimates for each covariate associated with the voters' characteristics.¹⁷

As shown in Table 7, the largest cross-quantile variation appears for the covariates *Abortion*, *Democrat*, *Republican*, and *Respondents' education*. Everything else being equal, preference heterogeneity is low for those who identify themselves as Democrats when voting for Clinton. The same is true for those who identify as Republicans when voting for Bush, which is in line with the literature on party alignment (see, e.g., Miller, 1991). However, when considering candidates from other parties, there is a considerable preference heterogeneity among voters. For example, in terms of abortion – a contentious issue during the election – pro-abortion voters have more coherent preferences when voting for Clinton rather than for Bush. This makes sense given the clear opposition of pro-abortion voters to Bush. Finally, even though educated voters tend to vote for Bush, they show higher heterogeneity in voting for Bush than in voting for Clinton (see also Lu, 2020). These diverse characteristics uncovered by examining conditional quantiles offer a richer understanding of the voting patterns among distinct voter groups, which are overlooked by models relying on conditional means.

¹⁷ The choice alternative Perot is set as the reference category as in the original analysis, and therefore has no estimates.

To assess the estimated coefficients for each variable, we can generate quantile coefficient plots for every choice alternative (Bush and Clinton), excluding the reference alternative (Perot). The task can be accomplished with the following straightforward line of code.

```
1 # Coefficient plot
2 plot(mod_vote1, "coef")
```

As Figure 16 shows, for the vote choice Bush, there is notable variation in the effects of most covariates across the distribution of the latent choice variable, i.e., the latent propensity of voting for Bush. Similarly, we can also plot the quantile coefficient values for vote choice Clinton, which is shown in Figure 17. This figure also shows a large variation in quantile effects of each covariate, which indicates significant heterogeneity in voting behaviors among American voters with diverse social, economic, and political backgrounds.

It is possible that unobserved heterogeneity among voters may affect the outcome of the vote when the ideological placement of the candidates changes. To draw a substantive interpretation of the estimation results, we can further compare the predicted vote shares of the three candidates by varying Bush's position while keeping the other variables constant (Alvarez & Nagler, 1995). It also enables us to evaluate how the quantile and mean-based models perform when compared with the real outcomes.

Figure 18 shows the predicted vote shares of the three candidates by varying Bush's position while keeping the other variables constant, with the solid lines representing the predictions of the multinomial probit model, and the dashed lines representing those of the CBQ model. The dots in the plot illustrate the actual vote share in the sample of the study. While the overall predictions are similar, the plot reveals that due to voter heterogeneity, the effects of the candidates' ideological movement vary. The largest variation occurs when voters decide to vote for or against Perot. By comparing the predicted vote shares and the real vote shares, it is clear that the multinomial probit estimator does not well account for unobserved heterogeneity among voters. The multinomial probit model is among the worst predictors when Bush's position is set at the actual value: 5.32. The best predictive quantile is 0.9, which suggests that loyal voters, who have a latent high propensity of voting for the corresponding candidate, have a greater influence on the final result than the rest of the voters. This finding suggests that, contrary to the conventional view emphasizing the importance of swing voters, the shift among core voters may have played a more significant role in determining the final result of the 1992 US presidential election. Thus, the underestimation of the multinomial model is likely due to




Figure 16 Estimated coefficients at quantiles ranging from 0.1 to 0.9 for US presidential election data with choice alternative Bush

Note: The estimates of variable *Ideological Distance* are the same between Bush and Clinton.

the fact that it only estimates the average preferences and does not take into account the variation of the preferences between different types of voters.

Taken together, the 1992 US election example illustrates how CBQ models can recover heterogeneity in multi-candidate choice settings that remains hidden in conventional multinomial models. By tracing covariate effects across




Figure 17 Estimated coefficients at quantiles ranging from 0.1 to 0.9 for US presidential election data with choice alternative Clinton

Note: The estimates of variable *Ideological Distance* are the same between Bush and Clinton.

the latent support distribution, the model not only improves predictive performance, but also yields substantively new insights into variation across types of voters. In this sense, the CBQ framework does more than refine existing estimates: it opens a different perspective on electoral behavior, one in which heterogeneous latent propensities matter directly for both explanation and prediction.




Figure 18 Predicted vote shares of three candidates by varying Bush's position

Note: The solid lines represent the predictions by the multinomial probit model, and the dashed lines represent the predictions by the quantile model.

4.4 Example of Varying Choice Alternatives: Government Formation

Compared to the previous application of vote choices, the data structure of government formation in multi-party parliamentary democracies is considerably more complex. The number of feasible coalition alternatives varies across countries and legislative terms, and the composition of those alternatives differs sharply from one formation opportunity to another. For instance, while the United Kingdom often presents only a small number of plausible governing combinations, postwar coalition bargaining in systems such as Germany involves many more viable party constellations. To investigate the mechanisms of government formation, Martin and Stevenson (2001) have generated one of the most comprehensive government formation datasets, which contains a total of 220 formation opportunities and 33,256 potential governments in 14 parliamentary democracies between 1945 and 1987.

In the analysis of government formation, the numbers of coalition alternatives differ across countries and elections, and the characteristics of potential governments within each formation opportunity can be correlated. For example, the status of the minority coalition and the minimal winning coalition may

depend on the inclusion or exclusion of the largest party in the coalition. Also, the coalition with the previous prime minister can be correlated with the status of the incumbent coalition. The IIA assumption is likely to be violated when unobserved government characteristics are correlated. This can happen when some government alternatives are substitutes or complements to each other. With a growing number of government alternatives, the risk of such correlation will also increase. Therefore, when estimating the traditional discrete choice model with equal weights over all coalition alternatives, a formation opportunity with a large number of coalition alternatives can distort the estimated formation likelihood of other coalitions.

To address variation in the number of coalition alternatives across cases, Martin and Stevenson (2001) propose to use the CL model, which accounts for varying formation opportunities across choice sets. However, the CL model relies on the strong IIA assumption, which is often violated in practice. In a later replication study, Glasgow, Golder, and Golder (2012) apply an MXL model to relax the restrictive IIA assumption and identify some disparities between their result and that of Martin and Stevenson (2001). Even so, both CL and MXL models assume a logistic form of the underlying error distribution and are thus restricted in assessing the full conditional distribution of the response variable (see the discussion in Section 3.2). They are therefore limited in their ability to reveal heterogeneity across the latent formation-propensity distribution. This is the setting in which the CBQ model becomes useful.

In the CBQ framework, conditional quantiles correspond to different points in the latent propensity distribution for successful coalition formation. The model therefore allows us to ask whether covariates matter differently for coalitions with relatively low, medium, or high latent probabilities of entering government. Although the underlying mathematics is more involved, estimation in practice is straightforward with the available software. The only additional requirement is that the model formula must include a choice-set identifier so that varying coalition alternatives are correctly grouped within each formation opportunity.

For a subset of the coalition formation data, the following codes estimate the 0.1th and 0.9th conditional quantiles while controlling for varying choice alternatives with a simple specification of the choice set indicator “`. | case`” in the `formula` object.¹⁸ Once estimation is complete, we can examine the estimates and their 95% credible intervals using the `print` function.

¹⁸ The correspondence between the variable names and their meanings is as follows: 1. `realg`: Really formed government; 2. `minor`: Minority Coalition; 3. `Minwin`: Minimal Winning Coalition; 4. `numpar`: Number of Parties in the Coalition; 5. `dompar`: Largest Party in the

```

1  # Take a subset of data_coalition with only the
    first 10 choice sets.
2  obs <- unique(data_coalition$case)[1:10]
3  data <- subset(data_coalition, data_coalition$
    case %in% obs)
4  ## Estimate the model
5  # 0.1th quantile
6  mod_coal1 <- cbq(realg ~ minor+ minwin + numpar +
    dompar + median + gdiv1 + mgodiv1 + prevpm +
    sq + mginvest + anmax2 + pro + anti|case, data
    = data, q = 0.1)
7  # 0.9th quantile
8  mod_coa9 <- cbq(realg ~ minor+ minwin + numpar +
    dompar + median + gdiv1 + mgodiv1 + prevpm +
    sq + mginvest + anmax2 + pro + anti|case, data
    = data, q = 0.9)
9  # Print the results
10 print(mod_coal1)
11 print(mod_coa9)

```

Based on the above estimation procedure, we can easily increase the number of estimated quantiles. Figure 19 shows the estimates of each covariate from 0.1th to 0.9th quantile. According to the figure, we can observe that the effects of most variables vary across quantiles. For example, while the effect of minority coalition remains significantly negative across quantiles, the magnitude decreases with increasing quantiles. Similarly, the positive effect of the largest party in the coalition decreases when a potential coalition becomes increasingly likely to be formed. These patterns suggest that average estimates obscure meaningful heterogeneity across different types of coalition alternatives.

Since most potential governments are not actually formed, there is a potential risk that the prediction accuracy can be inflated when all potential governments are included in the estimation. With the increasing number of choice alternatives, the prediction accuracy will automatically increase, although nothing else has been changed. This can be problematic when we want to compare the actual performance of different models. Indeed, the prediction accuracies of all models are close to 99.2% when all potential coalition governments are considered.

Coalition; 6. median: Median Party in the Coalition; 7. gdiv1: Ideological Divisions in the Coalition; 8. mgodiv1: Ideological Divisions within Majority Opposition; 9. prevpm: Previous Prime Minister in the Coalition; 10. sq: Incumbent Coalition; 11. mginvest: Minority Coalition where Investiture Vote Required; 12. anmax2: Anti-System Presence in the Coalition; 13. pro: Pre-Electoral Pact associated with the Coalition; 14. anti: Anti-Pact associated with the Coalition; 15. case: choice set indicator (Martin & Stevenson, 2001).




Figure 19 Estimated coefficients at quantiles ranging from 0.1 to 0.9 for government formation data

To address this problem, instead of calculating the prediction accuracy of all potential coalitions, we can compare the predicted probabilities and the prediction accuracy of those governments that were actually formed (Lu, 2020). For the 220 actually formed governments, the prediction accuracy of the CL and MXL models is 40.5% and 40%, respectively. The prediction accuracy of the CBQ model ranges from 34.5% to 42.3%. The best predictive quantiles are 0.1 and 0.9, with prediction accuracy of 42.3% and 40.9%, and the worst

predictive quantiles are in the middle. This indicates that models centered on the middle of the latent distribution do not summarize all coalition types equally well, which further justifies the use of quantile models. Overall, compared to the quantile models, conditional mean-based models such as CL and MXL are restrictive in their capacities to capture unobserved heterogeneity of potentially promising (high quantiles) and very unpromising coalition governments (low quantiles).

To draw a substantive interpretation of the results, we can further perform counterfactual analysis by setting substantively meaningful scenarios. However, due to the complex data structure of varying choice alternatives, the traditional counterfactual analysis cannot be easily applied in the context of coalition government formation. As Glasgow, Golder, and Golder (2012) have pointed out, varying a single variable while keeping the others constant is likely to produce illogical results. Following the advice, we can create logically coherent counterfactual scenarios by switching the largest party to the second-largest party, which consequently leads to changes in the features of other potential governments (Lu, 2020). The predicted probabilities of each model are then calculated from this logically coherent counterfactual dataset.

Using the full sample, Figure 20 plots the substitution patterns and compares changes in the predicted probabilities between CBQ and CL and between CBQ and MXL (see also figure 5 in Lu, 2020). The upper panels show the comparison between CBQ and CL, and the lower panels show the comparison between CBQ and MXL. The hollow points represent pairs of changes in the predicted probabilities when the largest party becomes the second-largest party. Black points are pairs whose differences are statistically significant at the 95% confidence level, and gray points are those that are insignificant. The dashed lines are 45-degree lines on which the changes in probabilities are the same.

It is clear from the figure that the patterns between the upper and lower panels are very similar and that the main difference between the quantile models and the CL/MXL model comes from the lower (0.1th) and upper (0.9th) quantiles. For those significantly different pairs, the lower quantile predicts, on average, a negative change in probabilities, while CL and MXL predict almost no change. In contrast, the upper quantile predicts no change in probabilities, while CL and MXL predict, on average, a negative change. The different substitution patterns between the lower and upper quantiles indicate that potentially promising coalitions (those in the upper quantiles) and misery coalitions (those in the lower quantiles) respond differently than an average government when the largest party becomes the second-largest party. In particular, the




Figure 20 Comparing substitution patterns of CBQ, CL, and MXL models (see also Figure 5 in Lu, 2020)

Note: The dashed lines are the 45-degree equal-division lines. Black points represent the pairs that are significantly different at a 95% confidence level while the gray points are insignificant. Q1 and Q9 indicate 0.1th and 0.9th quantile estimators, respectively. The average difference is calculated based on significantly different pairs.

promising coalitions suffer little from losing the largest party, while the misery coalitions' chances of entering government decrease. However, this formation heterogeneity among different types of coalitions has not been captured by the conditional mean-based CL or MXL models.

To further account for potential confounders, we may also want to incorporate mixed effects (fixed and/or random effects) into the estimation of the CBQ model. To do so, we can modify the syntax to “ $y \sim x \mid \text{indicator} \mid \text{variable of fixed effects} \mid \text{variable of random effects}$.” For example, if we are interested in estimating the random effects, the estimation formula can be specified as “ $y \sim x \mid \text{indicator} \mid 1 \mid \text{variable of random effects}$.” For the coalition formation data, we can further incorporate country-fixed effects with two explanatory variables as follows.

```

1 # Take a subset of data_coalition
2 obs <- unique(data_coalition$case)[c(1,40,50)]
3 data <- subset(data_coalition, data_coalition$
   case %in% obs)
4 # Estimate the model
5 model4 <- cbq(realg ~ minor + minwin|case|country
   |1, data = data, q = 0.5)

```

Taken together, the coalition-formation example shows that mean-based choice models provide only a partial account of coalition bargaining when substantial unobserved heterogeneity is present across potential governments. The CBQ estimates reveal that many coalition characteristics have markedly different effects at different points of the latent formation-propensity distribution. The counterfactual analysis reinforces this conclusion: relatively promising and highly unlikely coalitions respond differently to changes in party composition, and these differences are obscured in mean-based models such as CL and MXL. In this setting, the quantile perspective does not merely refine existing estimates—it changes the substantive picture of how coalition alternatives compete to form governments.

4.5 Summary

This section has extended the quantile framework to discrete-choice settings with multiple alternatives and alternative-specific features. In particular, it introduces the CBQ model for assessing potential diverse effects at various points along the conditional response distribution when there are multiple alternatives in choice sets.

Examples from the US presidential election and coalition formation in parliamentary democracies are used to demonstrate the potential of quantile models in generating new theoretical and empirical insights. In the electoral example, the CBQ model reveals substantial heterogeneity across the latent support distribution and suggests that mean-based models miss important differences in how voters respond to candidates and issues. This heterogeneity is reflected not only in the estimated coefficients, but also in the improved predictive performance.

The coalition-formation example points to a similarly important form of heterogeneity. Coalitions with relatively high latent propensities of formation appear to respond differently to changes in party composition than coalitions in the lower tail of the distribution. In particular, the results suggest that highly promising coalitions are less affected by the loss of largest-party status than highly unlikely coalitions. Such differences are obscured by mean-based choice models, but become visible once the quantile perspective is introduced.

Overall, these examples show that CBQ models do more than refine estimation in multiple-choice settings. They open up new ways of analyzing political choice by allowing researchers to study how effects vary across the latent distribution of support, viability, or selection. The accompanying code and software implementations can be adapted to a wide range of applied research settings.

5 A Brief Overview of Other Quantile Methods

5.1 Introduction

It is straightforward to expand the quantile modeling approach for the analysis of political heterogeneity in research situations other than what has been introduced in the previous chapters. To demonstrate the broader applicability of quantile models besides the examples presented in the previous chapters, this section will explore briefly, but not exhaustively, some additional quantile techniques that could be helpful for political research. These include in particular quantile models for estimating quantile treatment effects and survival data. While they are relatively less frequently used than the quantile models introduced in the previous chapters, these quantile models can be helpful when researchers attempt to identify causal relationships from experiments, or study different types of response variables beyond those discussed so far.

In this section I will also introduce variable selection and model comparison in quantile settings. In addition to the benefits of quantile models, this section will discuss in the concluding section the practical suggestions for using the quantile approach, such as those related to the quantile interpretation, the sample size, and the selection of the number of quantiles.

Although this is by no means a complete overview of the rapidly developing field of quantile methods, this section is intended to demonstrate the wide applicability of quantile models to broader application situations of political research. For other quantile methods such as those concerning non-parametric quantile regression and quantile estimation of longitudinal and time series data that have been even less frequently used in political science, interested readers may refer to Koenker (2017), Uribe and Guillen (2020), and Di Marzio, Panzera, and Taylor (2016), among others, for further details of implementation.

5.2 Quantile Treatment Effects

Quantile treatment effects (QTEs) describe how the effect of a treatment varies across the outcome distribution (Frölich & Melly, 2013). In the potential-outcomes framework, the τ th QTE is defined as the difference between the

τ th quantile of the treated outcome distribution and the τ th quantile of the untreated outcome distribution. In the simple case of a binary treatment T and a continuous outcome Y , this can be written intuitively as:

$$\text{QTE} = Q_{\tau}(Y|T = 1) - Q_{\tau}(Y|T = 0). \quad (5.1)$$

When a set of covariates X can also be controlled for, one may consider conditional quantile treatment effects of the form

$$\text{QTE} = Q_{\tau}(Y|X, T = 1) - Q_{\tau}(Y|X, T = 0). \quad (5.2)$$

In practice, the quantile treatment effects can be estimated by fitting quantile models for both treatment and control groups, and then comparing the estimated quantile-specific coefficients. This approach provides insights into how the treatment effect varies across different segments of the population, and thereby offers a more comprehensive understanding of the causal relations. The formula for estimating quantile treatment effects in the context of quantile regression can be expressed as follows:

$$Q_{\tau}(Y_i|X_i, T_i) = X_i'\beta_{\tau} + T_i\delta_{\tau} \quad (5.3)$$

where T_i is the treatment indicator variable, taking the value of 1 if the individual received treatment and 0 otherwise, X is a set of controls, and δ_{τ} captures the treatment contrast at quantile τ . The model assumes that the effect of treatment (T_i) on the conditional quantile of Y_i varies across different quantiles. Thereby, it allows for a more nuanced understanding of how the treatment effect changes at different points of the response distribution.

Estimating the quantile-specific coefficients and treatment effects follows the optimization procedure introduced in the previous chapters. Similar to the estimation procedure for observational studies, there are hands-on functions to estimate the quantile treatment effects when the data come from randomized experiments. The following example illustrates a slightly different but closely related use of quantile methods in experimental research: rather than estimating a treatment effect on the main substantive outcome, it uses quantile analysis to examine whether a key identification assumption holds uniformly across the response distribution.

As a real-world political example, consider a case study on vote buying in Nicaragua (Gonzalez-Ocantos et al., 2012). Vote buying, or more generally political clientelism, is a pervasive phenomenon in the politics of the developing world (Kitschelt, 2000). However, due to potential social desirability and concerns about legal repercussions, many recipients of political gifts may not truthfully disclose their behavior. To mitigate response bias, Gonzalez-Ocantos et al. (2012) employed an indirect measurement technique

called the “list experiment”. Rather than directly asking the respondents about sensitive issues like vote buying, this approach asks them to provide aggregate responses to a set of survey items that include the sensitive question (Blair & Imai, 2012). However, using a list experiment comes at the cost of reduced estimation efficiency, limiting researchers to estimating only the prevalence rather than individual instances of sensitive behaviors from the data (Blair, Coppock, & Moor, 2020; Glynn, 2013).

To enhance the efficiency of estimation in list experiments, Aronow et al. (2015) propose to combine direct questions with list experiments, which significantly reduces the uncertainty in estimating behaviors such as vote buying (Lu & Traummüller, 2026). However, ensuring accurate estimation through this method relies on the treatment independence assumption. This assumption posits that the assignment of treatment (i.e., inclusion in the list experiment treatment group) is independent of respondents’ answers to direct questions. In addition to placebo tests introduced by Aronow et al. (2015), we can examine the treatment independence assumption through quantile analysis between treatment assignment and responses to direct questions.

The rationale behind conducting quantile tests for the treatment independence assumption stems from the possibility of varying relationships between direct questions and treatment assignment across different segments of respondents. For example, individuals less likely to receive the treatment might also exhibit lower propensity to affirm the sensitive question in the direct form. Therefore, even if, on average, there is no discernible treatment effect on direct responses, the assumption of treatment independence could be compromised by significant effects in scenarios where treatment assignment probabilities differ. To explore potential heterogeneity of the treatment effect on direct responses, we can employ the quantile model by which the direct question is regressed against the treatment assignment. By estimating the relationship at different points along the distribution (quantiles), we gain insights into whether the assumption of treatment independence holds uniformly or if there are nuanced dependencies across different levels of the direct question’s response distribution. This approach allows us to detect any disparities in treatment effects that may exist among subgroups defined by their likelihood of affirming the sensitive question directly.

For the vote-buying example, Gonzalez-Ocantos et al. (2012) include in their list experiments also direct questions asking respondents whether they have received political gifts themselves (individual gifts) or are aware of some neighbors who have received gifts (neighbor gifts). Therefore, we can examine whether there are heterogeneous quantile treatment effects on the two direct questions. The following code presents the estimation procedure of both the

simple binary choice model and the quantile models. For simplicity, the code estimates the models separately at each quantile. In practice, this can also be automated in a loop.

```

1 library(readstata13)
2 library(cbq)
3 # Load Data
4 dat_vote <- read.dta13("voteBuying.dta")
5 # Baseline Logistic Model
6 mod_glm <- glm(neigift ~ treatment, data = dat_vote
  ,family = "binomial")
7 summary(mod_glm)
8 # Quantile Models for Neighbor Gifts
9 mod_cbq11 <- cbq(neigift ~ treatment, data = dat_
  vote, q = 0.1)
10 mod_cbq12 <- cbq(neigift ~ treatment, data = dat_
  vote, q = 0.2)
11 mod_cbq13 <- cbq(neigift ~ treatment, data = dat_
  vote, q = 0.3)
12 mod_cbq14 <- cbq(neigift ~ treatment, data = dat_
  vote, q = 0.4)
13 mod_cbq15 <- cbq(neigift ~ treatment, data = dat_
  vote, q = 0.5)
14 mod_cbq16 <- cbq(neigift ~ treatment, data = dat_
  vote, q = 0.6)
15 mod_cbq17 <- cbq(neigift ~ treatment, data = dat_
  vote, q = 0.7)
16 mod_cbq18 <- cbq(neigift ~ treatment, data = dat_
  vote, q = 0.8)
17 mod_cbq19 <- cbq(neigift ~ treatment, data = dat_
  vote, q = 0.9)
18 # Quantile Models for Individual Gifts
19 mod_cbq21 <- cbq(indgift ~ treatment, data = dat_
  vote, q = 0.1)
20 mod_cbq22 <- cbq(indgift ~ treatment, data = dat_
  vote, q = 0.2)
21 mod_cbq23 <- cbq(indgift ~ treatment, data = dat_
  vote, q = 0.3)
22 mod_cbq24 <- cbq(indgift ~ treatment, data = dat_
  vote, q = 0.4)
23 mod_cbq25 <- cbq(indgift ~ treatment, data = dat_
  vote, q = 0.5)
24 mod_cbq26 <- cbq(indgift ~ treatment, data = dat_
  vote, q = 0.6)
25 mod_cbq27 <- cbq(indgift ~ treatment, data = dat_
  vote, q = 0.7)
26 mod_cbq28 <- cbq(indgift ~ treatment, data = dat_
  vote, q = 0.8)
27 mod_cbq29 <- cbq(indgift ~ treatment, data = dat_
  vote, q = 0.9)

```

```

28 # Print the Estimates
29 print(mod_cbq11); print(mod_cbq12); print(mod_cbq13)
30 print(mod_cbq14); print(mod_cbq15); print(mod_cbq16)
31 print(mod_cbq17); print(mod_cbq18); print(mod_cbq19)
32 print(mod_cbq21); print(mod_cbq22); print(mod_cbq23)
33 print(mod_cbq24); print(mod_cbq25); print(mod_cbq26)
34 print(mod_cbq27); print(mod_cbq28); print(mod_cbq29)

```

Figure 21 shows the estimation results. For both the individual and neighbor gifts shown in the left and right panels of the figure, we find no significant treatment effects across all the specified quantiles from 0.1 to 0.9. This provides strong evidence that the treatment independence assumption is unlikely to be violated in this example and that direct questions can reasonably be combined with the list experiment, as long as the underlying bias of the direct questions is properly controlled for (Lu & Traunmüller, 2026).

The above example illustrates one simple application where the estimation of quantile treatment effects can help to reveal the distributional structure of the relationship between the treatment and response variable of interest. Under nonexperimental environments, quantile treatment effects may also be identified with other techniques such as the instrumental variable quantile regression (Abadie, Angrist, & Imbens, 1998; Chernozhukov & Hansen, 2005; Frölich & Melly, 2010). Interested readers may refer to the relevant packages, such as “IVQR” and their documentation, for further implementation details.




Figure 21 Estimates of quantile treatment effects

5.3 Quantile Model for Survival Data

Quantile models can also be applied to various types of response variables other than the ones illustrated in the previous chapters. One such response variable commonly encountered in political research is the time until a certain political event occurs, which is the subject of classical survival analysis. This section will thus demonstrate the use of quantile models for the analysis of survival data.

Quantile models for survival analysis extend the basic quantile framework to handle censored survival time data. Instead of modeling the hazard function, quantile models for survival analysis can be formulated as a generalization of the accelerated failure time model, which models directly the time until an event occurs.¹⁹ Thus, a positive coefficient means that an increase in the value of a covariate will increase the time until a certain event occurs. According to Peng and Huang (2008), given $Y = \log(T)$ and under certain conditions, we can formulate the quantile survival model as follows:

$$Q_{\tau}(T_i|X_i) = \exp \{X_i' \beta_{\tau}\} \quad (5.6)$$

The concrete estimation procedure will additionally account for the censoring probability of each observation. Compared to the conventional survival model, such as the Cox proportional hazards model, the quantile regression approach allows for the estimation of quantile-specific effects in survival analysis, thereby providing a more detailed understanding of how covariates influence different points of the survival time distribution.

To demonstrate the application of quantile models for survival analysis in political science, take political responses to the COVID-19 pandemic in Europe as an example (Toshkov, Carroll, & Yesilkagit, 2022). In response to the rapid spread of the COVID-19 pandemic, national governments across Europe had introduced various public policy measures such as lockdowns, school closures, and mask-wearing requirements. The timing of these measures varies across countries and over time, and may be further influenced by a number of political

¹⁹ In comparison to the accelerated failure time model, the Cox proportional hazards model adopts the following formula:

$$\lambda(t|X) = \lambda_0(t) \exp \{X' \beta\} \quad (5.4)$$

where $\lambda(t|X)$ is the conditional hazard function, and $\lambda_0(t)$ is the baseline hazard function. The unconditional hazard function evaluated at time t is defined as the instantaneous rate of occurrence:

$$\lambda(t) = \lim_{\Delta t \rightarrow 0} \frac{Pr(t \leq T < t + \Delta t | T \geq t)}{\Delta t} \quad (5.5)$$

where T is the true event time.

factors such as government capacity, political institutions, and party ideologies (Toshkov, Carroll, & Yesilkagit, 2022).

To examine what accounts for the diversity of policy responses to the pandemic, Toshkov, Carroll, and Yesilkagit (2022) collected a dataset on the announcement dates of anti-pandemic policies, including lockdowns and school closures, along with other explanatory variables such as government effectiveness, trust in government and type of government coalitions. Using their data, we attempt to examine in this example whether trust in government has varying effects on the timing of lockdowns depending on the relatively temporal location of each lockdown measure. Here, an event is an official announcement of the lockdown, and the response variable is the number of days between the first confirmed COVID-19 case in a country and the policy measure adoption (Toshkov, Carroll, & Yesilkagit, 2022).

As shown in Figure 22, with the elapsed time since the first confirmed COVID-19 case, the probability of no lockdowns decreases. In other words, the likelihood of imposing lockdowns after the confirmation of the first case increases over time. On average, the probability of introducing lockdown measures rises to 50% after 30 days since the first case, and it increases to almost 100% in 60 days. In fact, 23 out of 26 European countries included in the data had imposed lockdowns in 60 days since the first confirmed case. The estimation of the Cox proportional hazards model, which was adopted for the original analysis by Toshkov, Carroll, and Yesilkagit (2022), further shows that trust in government will slow down the introduction of lockdown measures. This may be due to the ease of sustaining compliance with social distancing measures in a high-trust environment so that the pressure for the government to introduce lockdowns is relatively low (Toshkov, Carroll, & Yesilkagit, 2022).

We can then proceed to analyze the potential heterogeneity in the effects of trust in government on the timing of lockdowns. Figure 23 shows the quantile estimation results for the survival data. Note that because the quantile models are performed on an accelerated failure time model (Portnoy, 2003) instead of the Cox proportional hazards model, positive coefficient values imply negative impact on hazard rates. In this example, a positive estimate means that an increase in government trust will postpone the introduction of lockdown measures. While the estimates from lower to middle quantiles (0.1th–0.4th quantiles) correspond to the results of the Cox proportional hazards model, we cannot find any significant effect of government trust on the timing of lockdown measures in the upper quantiles (≥ 0.5 th quantile).²⁰ This implies that

²⁰ The large variation in the estimates at the upper quantiles is partly due to the right-skewed distribution of the response variable.




Figure 22 Kaplan–Meier plot for introduction of lockdown measures




Figure 23 Coefficient estimates of quantile model for lockdown example

when the introduction of lockdown measures has been delayed too long, the public trust in government will no longer play a role in influencing the timing of policy introduction. These heterogeneous effects are likely due to the increasing risk of pandemic spread without lockdowns. As a result, over a certain threshold of the infection rate, the government is forced to impose lockdowns

regardless of the public trust in government, which explains the non-effect in the upper quantiles, where much time has passed without lockdowns since the first confirmed case. This is a substantively important pattern that would be obscured in a conventional hazard-based analysis.

5.4 Variable Selection and Model Comparison

In many empirical studies, researchers may want to identify the most relevant variables from a larger set of potential predictors. By selecting the most important variables, we can simplify and improve the accuracy and interpretability of quantile models. Variable selection helps to reduce computational complexity, identify key variables that influence the response, and may also improve the generalizability of the model to new data.

To perform variable selection in quantile settings, researchers have extended penalization methods to quantile regression building on the logic of LASSO (Least Absolute Shrinkage and Selection Operator) (Tibshirani, 1996). It is an extension of quantile regression that includes a penalty term to enforce sparsity in the parameter estimates. The formula for LASSO quantile regression is expressed as follows:

$$\text{minimize } \left\{ \sum_{i=1}^n \rho_{\tau}(y_i - \mathbf{x}_i' \boldsymbol{\beta}_{\tau}) + \lambda \sum_{j=1}^p |\beta_{\tau j}| \right\} \quad (5.7)$$

where λ is the regularization parameter that controls the strength of the penalty, and $|\beta_{\tau j}|$ represents the absolute value of the j -th coefficient in the vector $\boldsymbol{\beta}_{\tau}$. The first part of the formula, $\sum_{i=1}^n \rho_{\tau}(y_i - \mathbf{x}_i' \boldsymbol{\beta}_{\tau})$, is the quantile regression loss function, which measures the fit of the model to the data. The second part, $\lambda \sum_{j=1}^p |\beta_{\tau j}|$, is the LASSO penalty term that discourages large values of the coefficients. The objective is to find the values of $\boldsymbol{\beta}_{\tau}$ that minimize the combined loss and penalty. The regularization parameter λ controls the trade-off between fitting the data well and keeping the parameter estimates sparse.

The optimization problem is typically solved using optimization algorithms, such as coordinate descent or proximal gradient methods, to find the values of $\boldsymbol{\beta}_{\tau}$ that minimize the combined objective function. The `rq` function from the “quantreg” package allows for performing LASSO quantile model, where the `method = "lasso"` argument indicates the estimation of LASSO.

In the following example, we simulate a relatively homogeneous dataset with ten covariates, whose coefficients are $\boldsymbol{\beta} = \{2, 0, 2, 3, 2, 0, 1, 0.5, 0.5, 0\}$ for all values of τ . The codes estimating the LASSO quantile model are shown in the following.

```

1 # Quantile Lasso
2 # Generate synthetic data
3 set.seed(123)
4 n <- 1000 # Number of observations
5 X <- matrix(rnorm(n * 10), ncol = 10) # Ten
   independent variables
6 coefs <- c(2,0,2,3,2,0,1,0.5,0.5,0)
7 Y <- X%% matrix(coefs) + rnorm(n) # Response
   variable
8 # Combine data into a data frame
9 data <- data.frame(Y, X)
10 # Specify the quantile of interest
11 tau <- c(0.1,0.5,0.9)
12 # Perform LASSO quantile regression for variable
   selection
13 lasso_model <- rq(Y ~ X1 + X2 + X3 + X4 + X5 + X6
   + X7 + X8 + X9 + X10, data = data, tau = tau,
   method = "lasso")

```

The results in the following code show that the LASSO quantile models at three specified quantiles all correctly shrink the coefficients of the covariates (X2 and X6) without discernible effects. This estimation procedure can be easily adopted for other political research situations in which researchers only need to input the regular model specifications.

```

1 > round(coef(lasso_model),3)
2           tau= 0.1 tau= 0.5 tau= 0.9
3 (Intercept)   -1.203    0.000    1.282
4 X1             1.953    1.954    1.883
5 X2             0.094    0.000    0.042
6 X3             1.904    2.016    2.058
7 X4             3.095    3.035    2.995
8 X5             2.029    2.011    1.980
9 X6             0.000    0.000    0.000
10 X7            0.958    0.930    0.917
11 X8            0.510    0.529    0.400
12 X9            0.523    0.488    0.466
13 X10           0.000    0.016    0.025

```

In addition to variable selection, model comparison in quantile regression can be done using various criteria, such as Root Mean Squared Error (RMSE), Mean Absolute Error (MAE), Check Loss (CL, a.k.a. Quantile Loss or Pin-ball Loss), Akaike Information Criterion (AIC), and Bayesian Information Criterion (BIC), among others. RMSE is a widely used metric to evaluate the accuracy of regression models. It measures the square root of the average squared differences between predicted and observed values:

$$\text{RMSE} = \sqrt{\frac{1}{N} \sum_{i=1}^N (y_i - \hat{y}_i)^2}. \quad (5.8)$$

Similarly, MAE is a robust metric for evaluating regression model performance, which measures the average absolute differences between predicted and observed values:

$$\text{MAE} = \frac{1}{N} \sum_{i=1}^N |y_i - \hat{y}_i|. \quad (5.9)$$

Unlike RMSE or MAE, which focus on central tendencies, the check loss allows modeling of any quantile by asymmetrically weighting residuals:

$$\text{CL} = \frac{1}{N} \sum_{i=1}^N \rho_{\tau}(y_i - \hat{y}_i), \quad (5.10)$$

where $\rho_{\tau}(\cdot)$ is the check function as in Equation 2.4. In some settings, researchers also report information-criterion-style measures such as AIC or BIC. The AIC and BIC are measures of the relative quality of a statistical model for a given set of data. They penalize models based on the likelihood and the number of parameters, thereby balancing the trade-off between the goodness of fit and the complexity of the model (Akaike, 1987; Kass & Raftery, 1995; Schwarz, 1978). The difference between the two measures is that AIC imposes a fixed penalty independent of sample size, while BIC's penalty term grows with sample size:

$$\text{AIC} = -2 \ln(L) + 2k, \quad (5.11)$$

$$\text{BIC} = -2 \ln(L) + k \ln(N), \quad (5.12)$$

where L is the likelihood of the model, N is the sample size, and k is the number of estimated parameters. A lower AIC or BIC value indicates the better performance of a model. When comparing models, the model with the lowest AIC/BIC is generally considered the best, assuming that the models being compared are all based on the same data. Their use in quantile regression, however, is less straightforward than in standard likelihood-based models, and they are best treated as supplementary rather than primary criteria unless the underlying estimation framework provides a clear likelihood interpretation. In most applied quantile work, predictive criteria and quantile-specific loss functions are more directly informative.

Recall the study on the impact of economic crises on political representation introduced in Section 2. We can use the previously mentioned criteria to compare the performance of different quantile specifications. The estimation procedure is shown in the following codes:

```

1 # Model comparison
2 mod1 <- rq(coopscore_1 ~ gdppcgrowth + openness +
  population_log + elec + nparties + events +
  coopscore_mean_1 + factor(country) + factor(year
    ), tau = c(0.25,0.5,0.75), data = data)
3 summary(mod1)
4 # Predicted values
5 preds <- predict(mod1)
6 pred1 <- preds[,1]
7 pred2 <- preds[,2]
8 pred3 <- preds[,3]
9 # RMSE
10 rmse1 <- sqrt(mean((data$coopscore_1 - pred1)^2))
11 rmse2 <- sqrt(mean((data$coopscore_1 - pred2)^2))
12 rmse3 <- sqrt(mean((data$coopscore_1 - pred3)^2))
13 which.min(c(rmse1,rmse2,rmse3))
14 # MAE
15 mae1 <- mean(abs(data$coopscore_1 - pred1))
16 mae2 <- mean(abs(data$coopscore_1 - pred2))
17 mae3 <- mean(abs(data$coopscore_1 - pred3))
18 which.min(c(mae1,mae2,mae3))
19 # Check Loss
20 check_loss <- function(y, y_pred, tau) {
21   residuals <- y - y_pred
22   loss <- ifelse(residuals >= 0, tau * residuals, (
     tau - 1) * residuals)
23   mean(loss)
24 }
25 cl1 <- check_loss(data$coopscore_1, pred1,0.25)
26 cl2 <- check_loss(data$coopscore_1, pred2,0.5)
27 cl3 <- check_loss(data$coopscore_1, pred3,0.75)
28 which.min(c(cl1,cl2,cl3))
29 # AIC
30 aics <- AIC(mod1)
31 which.min(aics)
32 # BIC
33 bics <- AIC(mod1) - 2*dim(coef(mod1))[1] + dim(coef
  (mod1))[1]*log(dim(data)[1])
34 which.min(bics)

```

In this illustration, RMSE favors the 0.75 quantile specification, whereas MAE favors the median model. Most of the remaining criteria also point toward the median quantile as the strongest single specification. This divergence is not surprising. Different criteria emphasize different aspects of model performance: RMSE penalizes large prediction errors more heavily, MAE is more robust to outliers, and check loss is tailored to the quantile objective itself. When the values are close across quantiles, as they are here, the practical conclusion is not that one quantile definitively dominates the others, but

that several quantile specifications contribute useful information about the distribution.

For this reason, model-comparison statistics should be treated as helpful guides rather than as definitive arbiters of quantile choice. In substantive applications, it is usually more informative to estimate a grid of quantiles and to examine how the conclusions vary across the distribution than to rely on a single “best” quantile selected by one criterion alone. Model comparison should therefore complement, not replace, a broader distributional analysis.

5.5 Summary

This section has introduced several additional extensions of the quantile framework, including quantile models for estimating treatment effects and survival data. We have also discussed ways to select variables and compare models in quantile settings, thereby extending the practical toolkit introduced in the previous sections. However, despite the ability of quantile models to capture the conditional distribution of the response variable, there are some practical issues that are worth considering when applying quantile methods in empirical research.

Firstly, interpreting the estimated coefficients in quantile regression can be more challenging compared to ordinary least squares regression, especially when there are interactions between predictors. The coefficients in quantile regression represent the change in the specific quantile of the response variable associated with a one-unit change in the predictor variable, which may not always align with traditional interpretation. In order to better interpret the results for real-world applications, it is advisable to calculate substantive quantities of interest, such as predicted probabilities of occurrence.

Secondly, quantile models sometimes require a larger sample size compared to least squares regression, especially to obtain stable and accurate estimates in upper and lower quantiles. This is because quantile models estimate the regression coefficients for each quantile separately, and thus require a larger number of observations to achieve precise estimation, especially for extreme quantile locations. While estimation with more conditional quantiles provides a more complete picture of the conditional relationship between explanatory and response variables, the number of quantiles that can be effectively estimated is constrained by the sample size of the data. For example, in a data set with 100 observations, the estimates in the 0.015th conditional quantile may be the same as those in the 0.02th quantile as the two quantiles are likely to refer to the same observation. In such situations, to have a more comprehensive examination of the conditional relations between variables of interest, researchers may want to

increase the sample size by collecting more observations or widen the interval between quantiles to be estimated.

Nonetheless, these considerations do not diminish the value of quantile methods; rather, they clarify the conditions under which they are most informative. When used carefully, quantile models allow researchers to move beyond average effects, to examine heterogeneity across the conditional distribution, and to develop richer empirical accounts of political processes than mean-based approaches alone can provide.

6 Conclusion

Political science routinely studies heterogeneous actors, institutions, and contexts (De Marchi & Laver, 2023; Laver, 2020a, 2020b). Yet much empirical analysis still relies primarily on mean-based models, which often provide only a partial account of variation beyond the conditional mean. Quantile models offer a different perspective. By estimating relationships at different points of the conditional distribution, they make it possible to study heterogeneity directly rather than absorbing it into the error term. Because quantiles are rank-based and less sensitive to extreme observations than means, they also provide useful robustness advantages in many applied settings.

Despite the wide application of quantile models in many other scientific disciplines, they have remained comparatively underutilized in political science. One reason is that the discipline has lacked a systematic introduction connecting these methods to its own substantive questions, outcome types, and theoretical debates. To bridge the gap, this Element has introduced a set of quantile models for analyzing both continuous and discrete response variables. Besides the main quantile models, this Element also briefly introduces additional quantile methods to demonstrate the wide applicability of the method. In addition, the practical issues of quantile models, such as model comparison, variable selection, interpretation of the estimates, and the impact of sample size on the quality of quantile estimation, have also been discussed.

To make quantile models more accessible to political scientists, this Element provides a variety of real-world political examples, such as electoral choices, legislative decision-making, and party competition, to demonstrate the usefulness of the quantile approach in a wide field of political science. It reveals how existing conditional mean-based approaches can be distorted due to the presence of unobserved heterogeneity, and how results can vary across different parts of the distribution once quantile models are used to uncover heterogeneity. All examples are accompanied by hands-on computational codes which can be modified for other research purposes.

As concluding remarks, it will be helpful to highlight why and when quantile models should be used for application purposes. *Why should quantile models be used?* Restricting our attention to the average prevents us from raising new questions and answering those that have not been satisfactorily addressed. As noted by Mosteller and Tukey (1977, 266), “[w]hat the regression curve does is give a grand summary for the averages of the distributions corresponding to the set of x ’s. We could go further and compute several different regression curves corresponding to the various percentage points of the distributions and thus get a more complete picture of the set. Ordinarily this is not done, and so regression often gives a rather incomplete picture.”

Classical mean-based regression only provides insight into the expected values of the response variable given a set of covariates and may lead to incomplete or substantively misleading inferences, whereas the quantile approach allows for the examination of the entire response distribution. Quantile models estimate effects not only at the center but also at quantile-specific tails of the response distribution, thus providing a more comprehensive understanding of the relationship between explanatory and response variables, which would otherwise remain unknown. When the error term is not normally distributed and when the variance of the response variable changes across values of the independent variables (i.e., heteroscedasticity), quantile models can be more robust and more informative than least squares estimators. Multiple quantile specifications can be used to approximate the conditional distribution of the response variable without having to rely on restrictive distributional assumptions. Furthermore, the quantile models have a number of computational advantages, such as being robust to outliers, allowing for Bayesian inference in the estimation procedure, as well as providing well-quantified uncertainty without relying on asymptotic properties, making them an attractive option for applied political research.

When should quantile models be used? Since different mechanisms may apply to different subgroups of a population, quantile models are used to reveal heterogeneous effects at different conditional quantiles of the response distribution. This is helpful for evaluating varying policy interventions and political mechanisms across subpopulations, and summarizing extreme or peripheral cases in political science, such as radical parties, rare political events, and voters that are hardly represented by the average population. For example, the empirical analysis of vote choice discussed in Section 4 reveals that the effects of ideological distance on voting decisions may depend on the latent voting propensity of different voters. Compared to the average swing voters, the 1992 US presidential election exemplifies that core voters are indeed more responsive to the ideological gap between themselves and the candidates. By specifying a

number of conditional quantiles, it allows us to determine the magnitude of variation in the main effects of interest. In other words, when the research focus is not only on the average effect but also its distribution and variation over the whole range of the response variable, quantile models help to reveal varying effects and hidden heterogeneity that would otherwise be unexplored by mean-based models.

Moreover, when data contain outliers, extreme observations or are distributed asymmetrically, quantile models can be adopted as a robustness alternative to the mean-based models. Instead of computing complicated statistics for detecting outliers and normality, one can simply re-run the original analysis replacing the conditional mean with the conditional median, i.e., by applying a median regression model. If the normality assumption holds, the only loss from applying quantile models is a slight reduction in statistical efficiency.

Political analysis gains when we look beyond averages. Quantile models are not universal substitutes for mean-based methods, but they are indispensable whenever distributional heterogeneity is itself part of the substantive question. With increasingly accessible software and a growing methodological toolkit, they offer political scientists a practical and powerful way to ask new empirical questions and to produce more nuanced accounts of political life.

References

- Abadie, A., Angrist, J. D., & Imbens, G. W. (1998). Instrumental Variables Estimation of Quantile Treatment Effects. *NBER Technical Working Paper* (0229).
- Abramson, S. F., & Boix, C. (2019). Endogenous Parliaments: The Domestic and International Roots of Long-term Economic Growth and Executive Constraints in Europe. *International Organization*, 73(4), 793–837.
- Akaike, H. (1987). Factor Analysis and AIC. *Psychometrika*, 52, 317–332.
- Alvarez, R. M., & Glasgow, G. (1999). Two-Stage Estimation of Nonrecursive Choice Models. *Political Analysis*, 8(2), 147–165.
- Alvarez, R. M., & Nagler, J. (1995). Economics, Issues and the Perot Candidacy: Voter Choice in the 1992 Presidential Election. *American Journal of Political Science*, 39(3), 714–744.
- Appleton, S., Knight, J., Song, L., & Xia, Q. (2009). The Economics Of Communist Party Membership: The Curious Case of Rising Numbers and Wage Premium during China's Transition. *The Journal of Development Studies*, 45, 256–275.
- Arceneaux, K., & Nicholson, S. P. (2024). Anchoring Political Preferences: The Psychological Foundations of Status Quo Bias and the Boundaries of Elite Manipulation. *Political Behavior*, 46, 751–775.
- Aronow, P. M., Coppock, A., Crawford, F. W., & Green, D. P. (2015). Combining List Experiment and Direct Question Estimates of Sensitive Behavior Prevalence. *Journal of Survey Statistics and Methodology*, 3(1), 43–66.
- Arvate, P., Firpo, S., & Pieri, R. (2021). Can Women's Performance in Elections Determine the Engagement of Adolescent Girls in Politics? *European Journal of Political Economy*, 70, 102045, 1–19.
- Beck, N., Katz, J. N., & Tucker, R. (1998). Taking Time Seriously: Time-series-cross-section Analysis with a Binary Dependent Variable. *American Journal of Political Science*, 42(4), 1260–1288.
- Bellows, J., & Miguel, E. (2009). War and Local Collective Action in Sierra Leone. *Journal of Public Economics*, 93 (11-12), 1144–1157.
- Benoit, D. F., Alhamzawi, R., & Yu, K. (2013). Bayesian Lasso Binary Quantile Regression. *Computational Statistics*, 28, 2861–2873.
- Benoit, D. F., & Van den Poel, D. (2012). Binary Quantile Regression: A Bayesian Approach Based on the Asymmetric Laplace Distribution. *Journal of Applied Econometrics*, 27, 1174–1188.

- Benoit, D. F., & Van den Poel, D. (2017). bayesQR: A Bayesian Approach to Quantile Regression. *Journal of Statistical Software*, 76(7), 1–32. <https://doi.org/10.18637/jss.v076.i07>.
- Bertelli, A., & Richardson, L. E. (2008). Ideological Extremism and Electoral Design. Multimember versus Single Member Districts. *Public Choice*, 137, 347–368.
- Betz, T. (2017). Trading Interests: Domestic Institutions, International Negotiations, and the Politics of Trade. *The Journal of Politics*, 79(4), 1237–1252.
- Blair, G., Coppock, A., & Moor, M. (2020). When to Worry about Sensitivity Bias: A Social Reference Theory and Evidence from 30 Years of List Experiments. *American Political Science Review*, 114(4), 1297–1315.
- Blair, G., & Imai, K. (2012). Statistical Analysis of List Experiments. *Political Analysis*, 20(1), 47–77.
- Blattman, C. (2009). From Violence to Voting: War and Political Participation in Uganda. *American Political Science Review*, 103(2), 231–247.
- Blomberg, S. B., Broussard, N. H., & Hess, G. D. (2011). New Wine in Old Wineskins? Growth, Terrorism and the Resource Curse in Sub-saharan Africa. *European Journal of Political Economy*, 27, S50–S63.
- Braumoeller, B. F. (2006). Explaining Variance; or, Stuck in a Moment We Can't Get Out of. *Political Analysis*, 14(3), 268–290.
- Breunig, C. (2011). Reduction, Stasis, and Expansion of Budgets in Advanced Democracies. *Comparative Political Studies*, 44(8), 1060–1088.
- Breunig, C., & Jones, B. D. (2011). Stochastic Process Methods with an Application to Budgetary Data. *Political Analysis*, 19(1), 103–117.
- Breunig, C., & Koski, C. (2020). Topping Off and Bottoming Out: Setting Budget Priorities through Executive Power. *Policy Studies Journal*, 48(2), 342–366.
- Brunell, T. L., Koetzle, W., DiNardo, J., Grofman, B., & Feld, S. L. (1999). The R2= .93: Where Then Do They Differ? Comparing Liberal and Conservative Interest Group Ratings. *Legislative Studies Quarterly*, 24(1), 87–101.
- Buchinsky, M. (1998). Recent Advances in Quantile Regression Models: A Practical Guideline for Empirical Research. *Journal of Human Resources*, 33(1), 88–126.
- Bueno de Mesquita, B. (1981). *The War Trap*. Yale University Press.
- Caldara, D., & Iacoviello, M. (2022). Measuring Geopolitical Risk. *American Economic Review*, 112(4), 1194–1225.
- Carter, D. B., & Signorino, C. S. (2010). Back to the Future: Modeling Time Dependence in Binary Data. *Political Analysis*, 18(3), 271–292.
- Cheng, Y., Yang, L., & Deng, S. (2022). Nonprofit Density and Distributional Equity in Public Service Provision: Exploring Racial/Ethnic Disparities in

- Public Park Access across US Cities. *Public Administration Review*, 82(3), 473–486.
- Chernozhukov, V., & Hansen, C. (2005). An IV Model of Quantile Treatment Effects. *Econometrica*, 73(1), 245–261.
- Chowanietz, C. (2011). Rallying around the Flag or Railing against the Government? Political Parties' Reactions to Terrorist Acts. *Party Politics*, 17(5), 673–698.
- Cleophas, T. J., & Zwinderman, A. H. (2021). *Quantile Regression in Clinical Research: Complete Analysis for Data at a Loss of Homogeneity*. Springer.
- Davino, C., Furno, M., & Vistocco, D. (2014). *Quantile Regression: Theory and Applications*. John Wiley & Sons.
- De Luca, G., & Verpoorten, M. (2015). Civil War, Social Capital and Resilience in Uganda. *Oxford Economic Papers*, 67(3), 661–686.
- De Marchi, S., & Laver, M. (2023). *The Governance Cycle in Parliamentary Democracies*. Cambridge University Press.
- Di Marzio, M., Panzera, A., & Taylor, C. C. (2016). Nonparametric Circular Quantile Regression. *Journal of Statistical Planning and Inference*, 170, 1–14.
- Ding, P. (2025). *Linear Model and Extensions*. Chapman & Hall/CRC.
- Farrell, H., & Héritier, A. (2004). Interorganizational Negotiation and Intraorganizational Power in Shared Decision Making: Early Agreements under Codecision and Their Impact on the European Parliament and Council. *Comparative Political Studies*, 37(10), 1184–1212.
- Florios, K., & Skouras, S. (2008). Exact Computation of Max Weighted Score Estimators. *Journal of Econometrics*, 146(1), 86–91.
- Frandsen, B. R., Frölich, M., & Melly, B. (2012). Quantile Treatment Effects in the Regression Discontinuity Design. *Journal of Econometrics*, 168(2), 382–395.
- Frölich, M., & Melly, B. (2010). Estimation of Quantile Treatment Effects with Stata. *The Stata Journal*, 10(3), 423–457.
- Frölich, M., & Melly, B. (2013). Unconditional Quantile Treatment Effects Under Endogeneity. *Journal of Business & Economic Statistics*, 31(3), 346–357.
- Furno, M., & Vistocco, D. (2018). *Quantile Regression: Estimation and Simulation*. Wiley.
- Gilligan, M. J., Pasquale, B. J., & Samii, C. (2014). Civil War and Social Cohesion: Lab-in-the-field Evidence from Nepal. *American Journal of Political Science*, 58(3), 604–619.
- Glasgow, G. (2011). Introduction to the Virtual Issue: Recent Advances in Discrete Choice Methods in Political Science. *Political Analysis*, 19(1), 1–3.

- Glasgow, G. (2022). *Interpreting Discrete Choice Models*. Cambridge University Press.
- Glasgow, G., Golder, M., & Golder, S. N. (2012). New Empirical Strategies for the Study of Parliamentary Government Formation. *Political Analysis*, 20(2), 248–270.
- Glynn, A. N. (2013). What Can We Learn with Statistical Truth Serum? Design and Analysis of the List Experiment. *Public Opinion Quarterly*, 77(S1), 159–172.
- Gonzalez-Ocantos, E., De Jonge, C. K., Meléndez, C., Osorio, J., & Nickerson, D. W. (2012). Vote Buying and Social Desirability Bias: Experimental Evidence from Nicaragua. *American Journal of Political Science*, 56(1), 202–217.
- Grossman, G., Manekin, D., & Miodownik, D. (2015). The Political Legacies of Combat: Attitudes toward War and Peace among Israeli Ex-combatants. *International Organization*, 69(4), 981–1009.
- Guinjoan, M., & Rodon, T. (2021). Let's Party! The Impact of Local Festivities on the Incumbent's Electoral Support. *Local Government Studies*, 47(5), 712–734.
- Haan, P., Peichl, A., Schrenker, A., Weizsäcker, G., & Winter, J. (2022). Expectation Management of Policy Leaders: Evidence from COVID-19. *Journal of Public Economics*, 209, 104659, 1–26.
- Hadzic, D., Carlson, D., & Tavits, M. (2020). How Exposure to Violence Affects Ethnic Voting. *British Journal of Political Science*, 50(1), 345–362.
- Häge, F. M. (2013). Coalition Building and Consensus in the Council of the European Union. *British Journal of Political Science*, 43(3), 481–504.
- Hao, L., & Naiman, D. Q. (2007). *Quantile Regression*. SAGE.
- Hausman, J. A., & Wise, D. A. (1978). A Conditional Probit Model for Qualitative Choice: Discrete Decisions Recognizing Interdependence and Heterogeneous Preferences. *Econometrica*, 46(2), 403.
- Hensher, D. A., & Greene, W. H. (2003). The Mixed Logit Model: The State of Practice. *Transportation*, 30(2), 133–176.
- Hill, S. J., Hopkins, D. J., & Huber, G. A. (2021). Not by Turnout Alone: Measuring the Sources of Electoral Change, 2012 to 2016. *Science Advances*, 7(17), eabe3272, 1–10.
- Hoffman, M. D., & Gelman, A. (2014). The No-U-turn Sampler: Adaptively Setting Path Lengths in Hamiltonian Monte Carlo. *Journal of Machine Learning Research*, 15(1), 1593–1623.
- Holbein, J. (2016). Left Behind? Citizen Responsiveness to Government Performance Information. *American Political Science Review*, 110(2), 353–368.

- Horowitz, J. L. (1992). A Smoothed Maximum Score Estimator for the Binary Response Model. *Econometrica*, 60(3), 505–531.
- Indridason, I. H. (2008). Does Terrorism Influence Domestic Politics? Coalition Formation and Terrorist Incidents. *Journal of Peace Research*, 45(2), 241–259.
- Jetter, M., Agudelo, A. M., & Hassan, A. R. (2015). The Effect of Democracy on Corruption: Income is Key. *World Development*, 74, 286–304.
- Kaniovski, S., & Mueller, D. C. (2006). Community Size, Heterogeneity and Voter Turnouts. *Public Choice*, 129, 399–415.
- Kass, R. E., & Raftery, A. E. (1995). Bayes Factors. *Journal of the American Statistical Association*, 90(430), 773–795.
- Kitschelt, H. (2000). Linkages between Citizens and Politicians in Democratic Politics. *Comparative Political Studies*, 33(6-7), 845–879.
- Koenker, R. (2005). *Quantile Regression*. Cambridge University Press.
- Koenker, R. (2017). Quantile Regression: 40 Years On. *Annual Review of Economics*, 9, 155–176.
- Koenker, R. (2025). quantreg: Quantile regression [Computer software-manual]. Retrieved from <https://CRAN.R-project.org/package=quantreg> (R package version 6.1, <https://CRAN.R-project.org/package=quantreg>)
- Koenker, R., & Bassett, G. (1978). Regression Quantiles. *Econometrica*, 46(1), 33–50.
- Koenker, R., Chernozhukov, V., He, X., & Peng, L. (2017). *Handbook of Quantile Regression*. Chapman and Hall/CRC.
- Koenker, R., & Hallock, K. F. (2001). Quantile Regression. *Journal of Economic Perspectives*, 15(4), 143–156.
- Koenker, R., & Machado, J. A. F. (1999). Goodness of Fit and Related Inference Process for Quantile Regression. *Journal of the American Statistical Association*, 94(448), 1296–1310.
- Kordas, G. (2006). Smoothed Binary Regression Quantiles. *Journal of Applied Econometrics*, 21(3), 387–407.
- Kotz, S., Kozubowski, T. J., & Podgorski, K. (2001). *The Laplace Distribution and Generalizations: A Revisit with Applications to Communications, Economics, Engineering, and Finance*. Springer Science+Business Media, LLC.
- Kotz, S., Kozubowski, T. J., & Podgorski, K. (2003). An Asymmetric Multivariate Laplace Distribution. *Working Paper*, 1–26.
- Kozumi, H., & Kobayashi, G. (2011). Gibbs Sampling Methods for Bayesian Quantile Regression. *Journal of Statistical Computation and Simulation*, 81(11), 1565–1578.

- Krause, G. A., & Byers, J. S. (2022). Confirmation Dynamics: Differential Vetting in the Appointment of US Federal Agency Leaders. *The Journal of Politics*, 84(2), 1189–1201.
- Krause, G. A., & Zarit, M. (2022). The Retraction of Policy Benefits across US Federal Agencies: Programmatic Cutbacks and Executive Control of US Federal Grant Retrenchments. *Public Administration Review*, 82(4), 682–691.
- Laver, M. (2020a). *Agent-based Models of Polarization and Ethnocentrism*. Cambridge University Press.
- Laver, M. (2020b). *Agent-based Models of Social Life: Fundamentals*. Cambridge University Press.
- Lewis-Beck, C., & Lewis-Beck, M. (2015). *Applied Regression: An Introduction* (Vol. 22). Sage.
- Lijphart, A. (1996). The Puzzle of Indian Democracy: A Consociational Interpretation. *American Political Science Review*, 90(2), 258–268.
- Lu, X. (2020). Discrete Choice Data with Unobserved Heterogeneity: A Conditional Binary Quantile Model. *Political Analysis*, 28(2), 147–167.
- Lu, X., & Traummüller, R. (2026). Improving Studies of Sensitive Topics Using Prior Evidence: An Informative Bayesian Approach for List Experiments. *Political Science Research and Methods* First View, 1–17.
- Makowsky, M. D., & Miller, S. C. (2014). Education, Intelligence, and Attitude Extremity. *Public Opinion Quarterly*, 78(4), 832–858.
- Martin, L. W., & Stevenson, R. T. (2001). Government Formation in Parliamentary Democracies. *American Journal of Political Science*, 45(1), 33–50.
- McFadden, D. (1974). Conditional Logit Analysis of Qualitative Choice Behavior. In P. Zarembka (Ed.), *Frontiers in Econometrics* (pp. 105–142). Academic Press.
- McGrath, L. F. (2015). Estimating Onsets of Binary Events in Panel Data. *Political Analysis*, 23(4), 534–549.
- Miller, W. E. (1991). Party Identification, Realignment, and Party Voting: Back to the Basics. *American Political Science Review*, 85(2), 557–568.
- Miller, W. E., Kinder, D. R., & Rosenstone, S. J. (1993). The National Election Studies. 1993. American National Election Study, 1992: Pre-and Post-election Survey CPS Early Release Version, computer file. *Ann Arbor: University of Michigan, Center for Political Studies, and Inter-University Consortium for Political and Social Research*.
- Mosteller, F., & Tukey, J. W. (1977). *Data Analysis and Regression: A Second Course in Statistics*. Addison-Wesley.

- Neumayer, E., Plümper, T., & Barthel, F. (2014). The Political Economy of Natural Disaster Damage. *Global Environmental Change*, 24, 8–19.
- Oh, M.- S., Park, E. S., & So, B.- S. (2016). Bayesian Variable Selection in Binary Quantile Regression. *Statistics and Probability Letters*, 118, 177–181.
- Okada, K., & Samreth, S. (2012). The Effect of Foreign Aid on Corruption: A Quantile Regression Approach. *Economics Letters*, 115(2), 240–243.
- Peng, L., & Huang, Y. (2008). Survival Analysis with Quantile Regression Models. *Journal of the American Statistical Association*, 103(482), 637–649.
- Poole, K. T. (2000). Nonparametric Unfolding of Binary Choice Data. *Political Analysis*, 8(3), 211–237.
- Portnoy, S. (2003). Censored Regression Quantiles. *Journal of the American Statistical Association*, 98(464), 1001–1012.
- Poulos, J. (2019). Land Lotteries, Long-term Wealth, and Political Selection. *Public Choice*, 178, 217–230.
- R Core Team. (2021). R: A language and environment for statistical computing [Computer software manual]. Vienna, Austria. <https://www.R-project.org/>.
- Rainey, C. (2016). Dealing with Separation in Logistic Regression Models. *Political Analysis*, 24(3), 339–355.
- Rasmussen, A. (2011). Early Conclusion in Bicameral Bargaining: Evidence from the Co-decision Legislative Procedure of the European Union. *European Union Politics*, 12(1), 41–64.
- Reed, C., & Yu, K. (2009). A Partially Collapsed Gibbs Sampler for Bayesian Quantile Regression. *Technical Report, Department of Mathematical Sciences, Brunel University London*.
- Rohner, D., Thoenig, M., & Zilibotti, F. (2013a). Seeds of Distrust: Conflict in Uganda. *Journal of Economic Growth*, 18, 217–252.
- Rohner, D., Thoenig, M., & Zilibotti, F. (2013b). War Signals: A Theory of Trade, Trust, and Conflict. *Review of Economic Studies*, 80(3), 1114–1147.
- Rosenberg, A. S., Knappe, A. J., & Braumoeller, B. F. (2017). Unifying the Study of Asymmetric Hypotheses. *Political Analysis*, 25(3), 381–401.
- Sartori, A. E. (2003). An Estimator for Some Binary-Outcome Selection Models Without Exclusion Restrictions. *Political Analysis*, 11(2), 111–138.
- Schafer, J., & Holbein, J. B. (2020). When Time is of the Essence: A Natural Experiment on How Time Constraints Influence Elections. *The Journal of Politics*, 82(2), 418–432.
- Schwarz, G. (1978). Estimating the Dimension of a Model. *The Annals of Statistics*, 6(2), 461–464.
- Stasavage, D. (2002). Private Investment and Political Institutions. *Economics & Politics*, 14(1), 41–63.

- Stigler, S. M. (1984). Studies in the History of Probability and Statistics xl Boscovich, Simpson and a 1760 Manuscript Note on Fitting a Linear Relation. *Biometrika*, 71(3), 615–620.
- Tibshirani, R. (1996). Regression Shrinkage and Selection via the Lasso. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 58(1), 267–288.
- Toshkov, D., Carroll, B., & Yesilkagit, K. (2022). Government Capacity, Societal Trust or Party Preferences: What Accounts for the Variety of National Policy Responses to the Covid-19 Pandemic in Europe? *Journal of European Public Policy*, 29(7), 1009–1028.
- Trautmüller, R., Murr, A., & Gill, J. (2015). Modeling Latent Information in Voting Data with Dirichlet Process Priors. *Political Analysis*, 23(1), 1–20.
- Uribe, J. M., & Guillen, M. (2020). *Quantile Regression for Cross-sectional and Time Series Data: Applications in Energy Markets Using R*. Springer.
- Voors, M. J., Nillesen, E. E. M., Verwimp, P. et al. (2012). Violent Conflict and Behavior: A Field Experiment in Burundi. *American Economic Review*, 102(2), 941–964.
- Weschle, S. (2019). The Impact of Economic Crises on Political Representation in Public Communication: Evidence from the Eurozone. *British Journal of Political Science*, 49(3), 1097–1116.
- Wilkinson, S. (2006). *Votes and Violence: Electoral Competition and Ethnic Riots in India*. Cambridge University Press.
- Winship, C., & Mare, R. D. (1992). Models for Sample Selection Bias. *Annual Review of Sociology*, 18(1), 327–350.
- Wood, E. J. (2003). *Insurgent Collective Action and Civil War in El Salvador*. Cambridge University Press.
- Yu, K., & Moyeed, R. A. (2001). Bayesian Quantile Regression. *Statistics & Probability Letters*, 54, 437–447.
- Yu, K., & Zhang, J. (2005). A Three-parameter Asymmetric Laplace Distribution and Its Extension. *Communications in Statistics – Theory and Methods*, 34(9-10), 1867–1879.

Acknowledgments

This Element is a substantial extension of Chapter 2 of my doctoral dissertation and the paper of Lu (2020). It has benefited immensely from the insights, encouragement, and thoughtful critiques of many people who read and commented on all or parts of the manuscript. My gratitude extends to Simon Hug, Thomas König, Margit Tavits, Richard Traunmüller, Simon Weschle, and the anonymous reviewers for their helpful comments and suggestions, as well as to editor Michael Alvarez for endorsing this Element and providing invaluable guidance throughout the review and publication stages. I am deeply thankful to Junyan for her unwavering support. I am grateful to each of them for their time, generosity, and intellectual engagement. Their support has sharpened my arguments, clarified the interpretation of the results, and strengthened the overall structure of the work. Due to limitations on word count, it is regrettably not possible to acknowledge everyone by name, but I extend my heartfelt thanks to all who helped in the development of this Element.

I acknowledge the support by the Publication Fund of the School of International Studies, Peking University, and the Fundamental Research Funds for the Central Universities, Peking University.

Quantitative and Computational Methods for the Social Sciences

R. Michael Alvarez

California Institute of Technology

R. Michael Alvarez has taught at the California Institute of Technology his entire career, focusing on elections, voting behavior, election technology, and research methodologies. He has written or edited a number of books (recently, *Computational Social Science: Discovery and Prediction*, and *Evaluating Elections: A Handbook of Methods and Standards*) and numerous academic articles and reports.

Betsy Sinclair

Washington University in St. Louis

Betsy Sinclair is Professor and Chair of Political Science at WashU. Her research focuses on social influence and American political behavior. She is a fellow of the Society of Political Methodology and has served as an associate editor of *Political Analysis* and in leadership roles in The Society of Political Methodology and Visions in Political Methodology.

About the Series

The Elements Series Quantitative and Computational Methods for the Social Sciences contains short introductions and hands-on tutorials to innovative methodologies. These are often so new that they have no textbook treatment or no detailed treatment on how the method is used in practice. Among emerging areas of interest for social scientists, the series presents machine learning methods, the use of new technologies for the collection of data and new techniques for assessing causality with experimental and quasi-experimental data.

Quantitative and Computational Methods for the Social Sciences

Elements in the Series

Unsupervised Machine Learning for Clustering in Political and Social Research
Philip D. Waggoner

Using Shiny to Teach Econometric Models
Shawna K. Metzger

Modern Dimension Reduction
Philip D. Waggoner

Text Analysis in Python for Social Scientists: Prediction and Classification
Dirk Hovy

Interpreting Discrete Choice Models
Garrett Glasgow

Survival Analysis: A New Guide for Social Scientists
Alejandro Quiroz Flores

Adaptive Inventories: A Practical Guide for Applied Researchers
Jacob M. Montgomery and Erin L. Rossiter

A Practical Introduction to Regression Discontinuity Designs: Extensions
Matias D. Cattaneo, Nicolas Idrobo and Rocío Titiunik

Bayesian Social Science Statistics: From the Very Beginning
Jeff Gill and Le Bao

External Validity and Evidence Accumulation
Tara Slough and Scott A. Tyson

Bayesian Social Science Statistics: Getting Productive
Jeff Gill and Le Bao

Quantile Modeling for Political Research
Xiao Lu

A full series listing is available at: www.cambridge.org/QCMSS